



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/12, 15/53, C07K 16/40, 16/18, 14/47, C12N 9/02, A61K 38/17, 38/44, 48/00, G01N 33/50</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/28031</b> <b>(43) International Publication Date:</b> 18 May 2000 (18.05.00)
<b>(21) International Application Number:</b> PCT/US99/26592 <b>(22) International Filing Date:</b> 10 November 1999 (10.11.99)  <b>(30) Priority Data:</b> 60/107,911                      10 November 1998 (10.11.98)      US 60/149,332                      17 August 1999 (17.08.99)        US 60/151,242                      27 August 1999 (27.08.99)        US  <b>(71) Applicant:</b> EMORY UNIVERSITY [US/US]; 2009 Ridgewood Drive, Atlanta, GA 30322 (US).  <b>(72) Inventors:</b> LAMBETH, J., David; 461 Emory Drive, Decatur, GA 30207 (US). LASSEGUE, Bernard, P.; 564 Emory Oaks Way, Decatur, GA 30033 (US). GRIENDLING, Kathy, K.; 1819 Kanawha Trail, Stone Mountain, GA 30087 (US). ARNOLD, Rebecca, S.; Tucker, GA (US). GUANGJIE, Cheng; Lilburn, GA (US).  <b>(74) Agents:</b> ELSEVIER, Lisa, C. et al.; Jones & Askew, LLP, 2400 Monarch Tower, 3424 Peachtree Road, N.E., Atlanta, GA 30326 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> NOVEL MITOGENIC REGULATORS   <b>(57) Abstract</b> <p>The present invention relates to new genes encoding for the production of novel proteins involved in generation of reactive oxygen intermediates that affect cell division. The present invention also provides vectors containing these genes, cells transfected with these vectors, antibodies raised against these novel proteins, kits for detection, localization and measurement of these genes and proteins, and methods to determine the activity of drugs to affect the activity of the proteins of the present invention.</p>		



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



5

## NOVEL MITOGENIC REGULATORS

The U.S. Government has a paid-up license in this invention and the right in limited circumstances to require the patent owner to license others on reasonable terms as provided for by the terms of National Institutes of Health grants HL38206 and HL58000.

10

### TECHNICAL FIELD

The present invention relates to the field of normal and abnormal cell growth, in particular mitogenic regulation. The present invention provides the following: nucleotide sequences encoding for the production of enzymes that are mitogenic regulators; amino acid sequences of these enzymes; vectors containing these nucleotide sequences; methods for transfecting cells with vectors that produce these enzymes; transfected cells; methods for administering these transfected cells to animals to induce tumor formation; and antibodies to these enzymes that are useful for detecting and measuring levels of these enzymes, and for binding to cells possessing extracellular epitopes of these enzymes.

20

25

### BACKGROUND OF THE INVENTION

Reactive oxygen intermediates (ROI) are partial reduction products of oxygen: 1 electron reduces  $O_2$  to form superoxide ( $O_2^-$ ), and 2 electrons reduce  $O_2$  to form hydrogen peroxide ( $H_2O_2$ ). ROI are generated as a byproduct of aerobic metabolism and by toxicological mechanisms. There is growing evidence for regulated enzymatic generation of  $O_2^-$  and its conversion to  $H_2O_2$  in a variety of cells. The conversion of  $O_2^-$  to  $H_2O_2$  occurs spontaneously, but is markedly accelerated by superoxide dismutase (SOD). High levels of ROI are associated with damage to biomolecules such as DNA, biomembranes and proteins. Recent evidence indicates generation of ROI under

30

35



normal cellular conditions and points to signaling roles for  $O_2^-$  and  $H_2O_2$ .

Several biological systems generate reactive oxygen. Phagocytic cells such as neutrophils generate large quantities of ROI as part of their battery of bactericidal mechanisms. Exposure of neutrophils to bacteria or to various soluble mediators such as formyl-Met-Leu-Phe or phorbol esters activates a massive consumption of oxygen, termed the respiratory burst, to initially generate superoxide, with secondary generation of  $H_2O_2$ , HOCl and hydroxyl radical. The enzyme responsible for this oxygen consumption is the respiratory burst oxidase (nicotinamide adenine dinucleotide phosphate-reduced form (NADPH) oxidase).

There is growing evidence for the generation of ROI by non-phagocytic cells, particularly in situations related to cell proliferation. Significant generation of  $H_2O_2$ ,  $O_2^-$ , or both have been noted in some cell types. Fibroblasts and human endothelial cells show increased release of superoxide in response to cytokines such as interleukin-1 or tumor necrosis factor (TNF) (Meier et al. (1989) *Biochem J.* 263, 539-545.; Matsubara et al. (1986) *J. Immun.* 137, 3295-3298). Ras-transformed fibroblasts show increased superoxide release compared with control fibroblasts (Irani, et al. (1997) *Science* 275, 1649-1652). Rat vascular smooth muscle cells show increased  $H_2O_2$  release in response to PDGF (Sundaresan et al. (1995) *Science* 270, 296-299) and angiotensin II (Griendling et al. (1994) *Circ. Res.* 74, 1141-1148; Fukui et al. (1997) *Circ. Res.* 80, 45-51; Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321), and  $H_2O_2$  in these cells is associated with increased proliferation rate. The occurrence of ROI in a variety of cell types is summarized in Table 1 (adapted from Burdon, R. (1995) *Free Radical Biol. Med.* 18, 775-794).



Table 1

<u>Superoxide</u>	<u>Hydrogen Peroxide</u>
human fibroblasts	Balb/3T3 cells
human endothelial cells	rat pancreatic islet cells
human/rat smooth muscle cells	murine keratinocytes
human fat cells	rabbit chondrocytes
human osteocytes	human tumor cells
BHK-21 cells	fat cells, 3T3 L1 cells
human colonic epithelial cells	

ROI generated by the neutrophil have a cytotoxic function. While ROI are normally directed at the invading microbe, ROI can also induce tissue damage (e.g., in inflammatory conditions such as arthritis, shock, lung disease, and inflammatory bowel disease) or may be involved in tumor initiation or promotion, due to damaging effects on DNA. Nathan (Szatrowski et al. (1991) *Canc. Res.* 51, 794-798) proposed that the generation of ROI in tumor cells may contribute to the hypermutability seen in tumors, and may therefore contribute to tumor heterogeneity, invasion and metastasis.

In addition to cytotoxic and mutagenic roles, ROI have ideal properties as signal molecules: 1) they are generated in a controlled manner in response to upstream signals; 2) the signal can be terminated by rapid metabolism of  $O_2^-$  and  $H_2O_2$  by SOD and catalase/peroxidases; 3) they elicit downstream effects on target molecules, e.g., redox-sensitive regulatory proteins such as NF kappa B and AP-1 (Schreck et al. (1991) *EMBO J.* 10, 2247-2258; Schmidt et al. (1995) *Chemistry & Biology* 2, 13-22). Oxidants such as  $O_2^-$  and  $H_2O_2$  have a relatively well defined signaling role in bacteria, operating via the SoxI/II regulon to regulate transcription.

ROI appear to have a direct role in regulating cell division, and may function as mitogenic signals in pathological conditions related to growth. These conditions include cancer and cardiovascular disease.  $O_2^-$  is generated in endothelial cells



in response to cytokines, and might play a role in angiogenesis (Matsubara et al. (1986) *J. Immun.* 137, 3295-3298).  $O_2^-$  and  $H_2O_2$  are also proposed to function as "life-signals", preventing cells from undergoing apoptosis (Matsubara et al. (1986) *J. Immun.* 137, 3295-3298). As discussed above, many cells respond to growth factors (e.g., platelet derived growth factor (PDGF), epidermal derived growth factor (EGF), angiotensin II, and various cytokines) with both increased production of  $O_2^-$  / $H_2O_2$  and increased proliferation. Inhibition of ROI generation prevents the mitogenic response. Exposure to exogenously generated  $O_2^-$  and  $H_2O_2$  results in an increase in cell proliferation. A partial list of responsive cell types is shown below in Table 2 (adapted from Burdon, R. (1995) *Free Radical Biol. Med.* 18, 775-794).

Table 2

<u>Superoxide</u>	<u>Hydrogen peroxide</u>
human, hamster fibroblasts	mouse osteoblastic cells
Balb/3T3 cells	Balb/3T3 cells
human histiocytic leukemia	rat, hamster fibroblasts
mouse epidermal cells	human smooth muscle cells
rat colonic epithelial cells	rat vascular smooth muscle cells
rat vascular smooth muscle cells	

While non-transformed cells can respond to growth factors and cytokines with the production of ROI, tumor cells appear to produce ROI in an uncontrolled manner. A series of human tumor cells produced large amounts of hydrogen peroxide compared with non-tumor cells (Szatrowski et al. (1991) *Canc. Res.* 51, 794-798). Ras-transformed NIH 3T3 cells generated elevated amounts of superoxide, and inhibition of superoxide generation by several mechanisms resulted in a reversion to a "normal" growth phenotype.

$O_2^-$  has been implicated in maintenance of the transformed phenotype in cancer cells including melanoma, breast carcinoma, fibrosarcoma, and virally transformed tumor



cells. Decreased levels of the manganese form of SOD (MnSOD) have been measured in cancer cells and *in vitro*-transformed cell lines, predicting increased  $O_2^-$  levels (Burdon, R. (1995) *Free Radical Biol. Med.* 18, 775-794). MnSOD is encoded on chromosome 6q25 which is very often lost in melanoma. Overexpression of MnSOD in melanoma and other cancer cells (Church et al. (1993) *Proc. of Natl. Acad. Sci.* 90, 3113-3117; Fernandez-Pol et al. (1982) *Canc. Res.* 42, 609-617; Yan et al. (1996) *Canc. Res.* 56, 2864-2871) resulted in suppression of the transformed phenotype.

ROI are implicated in growth of vascular smooth muscle associated with hypertension, atherosclerosis, and restenosis after angioplasty.  $O_2^-$  generation is seen in rabbit aortic adventitia (Pagano et al. (1997) *Proc. Natl. Acad. Sci.* 94, 14483-14488). Vascular endothelial cells release  $O_2^-$  in response to cytokines (Matsubara et al. (1986) *J. Immun.* 137, 3295-3298).  $O_2^-$  is generated by aortic smooth muscle cells in culture, and increased  $O_2^-$  generation is stimulated by angiotensin II which also induces cell hypertrophy. In a rat model system, infusion of angiotensin II leads to hypertension as well as increased  $O_2^-$  generation in subsequently isolated aortic tissue (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321.; Yu et al. (1997) *J. Biol. Chem.* 272, 27288-27294). Intravenous infusion of a form of SOD that localizes to the vasculature or an infusion of an  $O_2^-$  scavenger prevented angiotensin II induced hypertension and inhibited ROI generation (Fukui et al. (1997) *Circ. Res.* 80, 45-51).

The neutrophil NADPH oxidase, also known as phagocyte respiratory burst oxidase, provides a paradigm for the study of the specialized enzymatic ROI-generating system. This extensively studied enzyme oxidizes NADPH and reduces oxygen to form  $O_2^-$ . NADPH oxidase consists of multiple proteins and is regulated by assembly of cytosolic and membrane components. The catalytic moiety consists of flavocytochrome  $b_{558}$ , an integral plasma membrane enzyme comprised of two components: gp91phox (gp refers to



glycoprotein; phox is an abbreviation of the words phagocyte and oxidase) and p22phox (p refers to protein). gp91phox contains 1 flavin adenine dinucleotide (FAD) and 2 hemes as well as the NADPH binding site. p22phox has a C-terminal  
5 proline-rich sequence which serves as a binding site for cytosolic regulatory proteins. The two cytochrome subunits, gp91phox and p22phox appear to stabilize one another, since the genetic absence of either subunit, as in the inherited disorder chronic granulomatous disease (CGD), results in the absence of  
10 the partner subunit (Yu et al. (1997) *J. Biol. Chem.* 272, 27288-27294). Essential cytosolic proteins include p47phox, p67phox and the small GTPase Rac, of which there are two isoforms. p47phox and p67phox both contain SH<sub>3</sub> regions and proline-rich regions which participate in protein interactions  
15 governing assembly of the oxidase components during activation. The neutrophil enzyme is regulated in response to bacterial phagocytosis or chemotactic signals by phosphorylation of p47phox, and perhaps other components, as well as by guanine nucleotide exchange to activate the GTP-binding protein Rac.  
20

The origin of ROI in non-phagocytic tissues is unproven, but the occurrence of phagocyte oxidase components has been evaluated in several systems by immunochemical methods, Northern blots and reverse transcriptase-polymerase  
25 chain reaction (RT-PCR). The message for p22phox is expressed widely, as is that for Rac1. Several cell types that are capable of O<sub>2</sub><sup>-</sup> generation have been demonstrated to contain all of the phox components including gp91phox, as summarized below in Table 3. These cell types include endothelial cells,  
30 aortic adventitia and lymphocytes.



Table 3

	Tissue	gp91phox	p22phox	p47phox	p67phox
	neutrophil	+ <sup>1,2</sup>	+ <sup>1,2</sup>	+ <sup>1,2</sup>	+ <sup>1,2</sup>
	aortic adventitia	+ <sup>1</sup>	+ <sup>1</sup>	+ <sup>1</sup>	+ <sup>1</sup>
5	lymphocytes	+ <sup>2</sup>	+ <sup>2</sup>	+ <sup>1,2</sup>	+ <sup>1,2</sup>
	endothelial cells	+ <sup>2</sup>	+ <sup>2</sup>	+ <sup>1,2</sup>	+ <sup>1,2</sup>
	glomerular mesangial cells	-	+ <sup>1,2</sup>	+ <sup>1,2</sup>	+ <sup>1,2</sup>
	fibroblasts	-	+ <sup>2</sup>	+ <sup>1,2</sup>	+ <sup>2</sup>
10	aortic sm. muscle	-	+ <sup>1,2</sup>	?	?

1= protein expression shown. 2= mRNA expression shown.

15 However, a distinctly different pattern is seen in several other cell types shown in Table 3 including glomerular mesangial cells, rat aortic smooth muscle and fibroblasts. In these cells, expression of gp91phox is absent while p22phox and in some cases cytosolic phox components have been demonstrated to be present. Since gp91phox and p22phox stabilize one another in the neutrophil, there has been much speculation that some molecule, possibly related to gp91phox, accounts for ROI generation in glomerular mesangial cells, rat aortic smooth muscle and fibroblasts (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321). Investigation of fibroblasts from a patient with a genetic absence of gp91phox provides proof that the gp91phox subunit is not involved in ROI generation in these cells (Emmendorffer et al. (1993) *Eur. J. Haematol.* 51, 223-227). Depletion of p22phox from vascular smooth muscle using an antisense approach indicated that this subunit participates in ROI generation in these cells, despite the absence of detectable gp91phox (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321). At this time the molecular candidates possibly related to gp91phox and involved in ROI generation in these cells are unknown.

35 Accordingly, what is needed is the identity of the proteins involved in ROI generation, especially in non-phagocytic tissues and cells. What is also needed are the



nucleotide sequences encoding for these proteins, and the primary sequences of the proteins themselves. Also needed are vectors designed to include nucleotides encoding for these proteins. Probes and PCR primers derived from the nucleotide sequence are needed to detect, localize and measure nucleotide sequences, including mRNA, involved in the synthesis of these proteins. In addition, what is needed is a means to transfect cells with these vectors. What is also needed are expression systems for production of these molecules. Also needed are antibodies directed against these molecules for a variety of uses including localization, detection, measurement and passive immunization.

#### SUMMARY OF THE INVENTION

The present invention solves the problems described above by providing a novel family of nucleotide sequences and proteins encoded by these nucleotide sequences termed mox proteins and duox proteins. In particular the present invention provides compositions comprising the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, and fragments thereof, which encode for the expression of proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, respectively, and fragments thereof. While not wanting to be bound by the following statement, it is believed that these proteins are involved in ROI production. The present invention also provides vectors containing these nucleotide sequences, cells transfected with these vectors which produce the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, and fragments thereof, and antibodies to these proteins and fragments thereof. The present invention also provides methods for stimulating cellular proliferation by administering vectors encoded for production of the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The present invention also



provides methods for stimulating cellular proliferation by administering the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The nucleotides and antibodies of the present invention are useful for the detection, localization and measurement of the nucleic acids encoding for the production of the proteins of the present invention, and also for the detection, localization and measurement of the proteins of the present invention. These nucleotides and antibodies can be combined with other reagents in kits for the purposes of detection, localization and measurement.

Most particularly, the present invention involves a method for regulation of cell division or cell proliferation by modifying the activity or expression of the proteins described as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof. These proteins, in their naturally occurring or expressed forms, are expected to be useful in drug development, for example for screening of chemical and drug libraries by observing inhibition of the activity of these enzymes. Such chemicals and drugs would likely be useful as treatments for cancer, prostatic hypertrophy, benign prostatic hypertrophy, hypertension, atherosclerosis and many other disorders involving abnormal cell growth or proliferation as described below. The entire expressed protein may be useful in these assays. Portions of the molecule which may be targets for inhibition or modification include but are not limited to the binding site for pyridine nucleotides (NADPH or NADH), the flavoprotein domain (approximately the C-terminal 265 amino acids), and/or the binding or catalytic site for flavin adenine dinucleotide (FAD).

The method of the present invention may be used for the development of drugs or other therapies for the treatment of conditions associated with abnormal growth including, but not limited to the following: cancer, psoriasis, prostatic hypertrophy, benign prostatic hypertrophy, cardiovascular disease, proliferation of vessels, including but



not limited to blood vessels and lymphatic vessels, arteriovenous malformation, vascular problems associated with the eye, atherosclerosis, hypertension, and restenosis following angioplasty. The enzymes of the present invention are excellent  
5 targets for the development of drugs and other agents which may modulate the activity of these enzymes. It is to be understood that modulation of activity may result in enhanced, diminished or absence of enzymatic activity. Modulation of the activity of these enzymes may be useful in treatment of  
10 conditions associated with abnormal growth.

Drugs which affect the activity of the enzymes represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, may also be combined with other therapeutics in the  
15 treatment of specific conditions. For example, these drugs may be combined with angiogenesis inhibitors in the treatment of cancer, with antihypertensives for the treatment of hypertension, and with cholesterol lowering drugs for the treatment of atherosclerosis.

Accordingly, an object of the present invention is to provide nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production.  
20

Another object of the present invention is to provide vectors containing these nucleotide sequences, or fragments thereof.  
25

Yet another object of the present invention is to provide cells transfected with these vectors.

Still another object of the present invention is to administer cells transfected with these vectors to animals and humans.  
30

Another object of the present invention is to provide proteins, or fragments thereof, that are involved in ROI production.  
35

Still another object of the present invention is to provide antibodies, including monoclonal and polyclonal



antibodies, or fragments thereof, raised against proteins, or fragments thereof, that are involved in ROI production.

Another object of the present invention is to administer genes containing nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans and also to cells obtained from animals and humans.

Another object of the present invention is to administer antisense complimentary sequences of genes containing nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans and also to cells obtained from animals and humans.

Yet another object of the present invention is to provide a method for stimulating or inhibiting cellular proliferation by administering vectors containing nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans. It is also an object of the present invention to provide a method for stimulating or inhibiting cellular proliferation by administering vectors containing antisense complimentary sequences of nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans. These methods of stimulating cellular proliferation are useful for a variety of purposes, including but not limited to, developing animal models of tumor formation, stimulating cellular proliferation of blood marrow cells following chemotherapy or radiation, or in cases of anemia.

Still another object of the present invention is to provide antibodies useful in immunotherapy against cancers expressing the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof.



Yet another object of the present invention is to provide nucleotide probes useful for the detection, localization and measurement of nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production.

Another object of the present invention is to provide antibodies useful for the detection, localization and measurement of nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production.

Another object of the present invention is to provide kits useful for detection of nucleic acids including the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins, or fragments thereof, that are involved in ROI production.

Yet another object of the present invention is to provide kits useful for detection and measurement of nucleic acids including the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins, or fragments thereof, that are involved in ROI production.

Still another object of the present invention is to provide kits useful for the localization of nucleic acids including the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins, or fragments thereof that are involved in ROI production.

Another object of the present invention is to provide kits useful for detection of proteins, including the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, that are involved in ROI production.

Yet another object of the present invention is to provide kits useful for detection and measurement of proteins,



including the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, that are involved in ROI production.

5 Still another object of the present invention is to provide kits useful for localization of proteins, including the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, that are involved in ROI production.

10 Yet another object of the present invention is to provides kits useful for the detection, measurement or localization of nucleic acids, or fragments thereof, encoding for proteins, or fragments thereof, that are involved in ROI production, for use in diagnosis and prognosis of abnormal cellular proliferation related to ROI production.

15 Another object of the present invention is to provides kits useful for the detection, measurement or localization of proteins, or fragments thereof, that are involved in ROI production, for use in diagnosis and prognosis of abnormal cellular proliferation related to ROI production.

20 These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiments and the appended drawings.

25

#### BRIEF DESCRIPTION OF THE FIGURES

Fig. 1(a-d). Comparison of amino acid sequences of the human mox1 protein (labeled mox1.human, SEQ ID NO:2), rat mox1 protein (labeled mox1.rat, SEQ ID NO:21), human mox2 protein (labeled mox2.human., SEQ ID NO:4) of the present invention to human (gp 91phox/human.pep, SEQ ID NO:12) bovine (gp 91 phox/bovine.pep, SEQ ID NO:37), and murine (gp 91 phox/mouse.pep, SEQ ID NO:38) proteins. Also included are related plant enzyme proteins cytb 558.arabidopsis.pep (SEQ ID NO:39) and cytb558.rice.pep,

30

35



(SEQ ID NO:40). Enclosed in boxes are similar amino acid residues.

5 Fig. 2. Sequence similarities among proteins related to gp91phox including human mox1 (SEQ ID NO:2), human mox2 (SEQ ID NO:4), and rat mox1 (SEQ ID NO:21). The dendrogram indicates the degree of similarity among this family of proteins, and also includes the related plant enzymes.

10 Fig. 3. Cell free assay for mox-1 activity. Superoxide generation was measured using the chemiluminescent reaction between lucigenin and superoxide in cell lysates from vector control NEF2 and mox1 transfected NIH3T3 cells.

15 Fig. 4. Superoxide generation by human mox1. Reduction of NBT in mox1 transfected and control fibroblasts was measured in the absence (filled bars) or presence (open bars) or superoxide dismutase.

20 Fig. 5. Aconitase (filled bars), lactate dehydrogenase (narrow hatching) and fumarase (broad hatching) were determined in lysates of cells transfected with vector-alone (NEF2) or with mox1 (YA26, YA28 and YA212).

#### DETAILED DESCRIPTION OF THE INVENTION

25 The present invention solves the problems described above by providing a novel family of nucleotide sequences and proteins, encoded by these nucleotide sequences, termed mox proteins and duox proteins. The term "mox" refers to "mitogenic oxidase" while the term "duox" refers to "dual oxidase". In particular, the present invention provides novel compositions comprising the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, and fragments thereof, which encode, respectively, for the expression of proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof.

35 Both the mox and duox proteins described herein have homology to the gp91phox protein involved in ROI



generation, however, the mox and duox proteins comprise a novel and distinct family of proteins. The mox proteins included in the present invention have a molecular weight of approximately 65 kDa as determined by reducing gel electrophoresis and are capable of inducing ROI generation in cells. As described in more detail below, the mox proteins of the present invention also function in the regulation of cell growth, and are therefore implicated in diseases involving abnormal cell growth such as cancer. The present invention describes mox proteins found in human and rat, however, it is likely that the mox family of genes/proteins is widely distributed among multicellular organisms.

The duox proteins described herein are larger than the mox proteins and have three distinct regions: the amino terminal region having homology to peroxidase proteins, the internal region having homology to calmodulin (CAM) proteins and the carboxy-terminal region having homology to mox proteins. Human duox1 is shown in SEQ ID NO:46 and a portion of human duox2 is shown in SEQ ID NO:48. Nucleotides encoding duox1 and duox2 proteins are also shown in SEQ ID NO: 45 and SEQ ID NO:47, respectively. In addition to the human duox proteins, comparison of the sequence of human duox1 and human duox2 with genomic databases using BLAST searching resulted in the identification of two homologs of duox in *C. elegans* (Ce-duox1 and Ce-duox2). *Drosophila* also appears to have at least one duox homolog. Thus, the duox family of genes/proteins is widely distributed among multicellular organisms.

Although not wanting to be bound by the following statement, it is believed that duox1 and duox2 have dual enzymatic functions, catalyzing both the generation of superoxide and peroxidative type reactions. The latter class of reactions utilize hydrogen peroxide as a substrate (and in some cases have been proposed to utilize superoxide as a substrate). Since hydrogen peroxide is generated spontaneously from the dismutation of superoxide, it is believed that the NAD(P)H



oxidase domain generates the superoxide and/or hydrogen peroxide which can then be used as a substrate for the peroxidase domain. In support of this hypothesis, a model for the duox1 protein in *C. elegans* has been developed that has an extracellular N-terminal peroxidase domain, a transmembrane region and a NADPH binding site located on the cytosolic face of the plasma membrane. By analogy with the neutrophil NADPH-oxidase which generates extracellular superoxide, human duox1 is predicted to generate superoxide and its byproduct hydrogen peroxide extracellularly where it can be utilized by the peroxidase domain.

While the ROI generated by duox1 and duox2 may function as does mox1 in regulation of cell growth, the presence of the peroxidase domain is likely to confer additional biological functions. Depending upon the co-substrate, peroxidases can participate in a variety of reactions including halogenation such as the generation of hypochlorous acid (HOCl) by myeloperoxidase and the iodination of tyrosine to form thyroxine by thyroid peroxidase. Peroxidases have also been documented to participate in the metabolism of polyunsaturated fatty acids, and in the chemical modification of tyrosine in collagen (by sea urchin ovoperoxidase). Although not wanting to be bound by this statement, it is believed that the predicted transmembrane nature of duox1 facilitates its function in the formation or modification of extracellular matrix or basement membrane. Since the extracellular matrix plays an important role in tumor cell growth, invasion and metastasis, it is believed that the duox type enzymes play a pathogenic role in such conditions.

In addition to the nucleotide sequences described above, the present invention also provides vectors containing these nucleotide sequences and fragments thereof, cells transfected with these vectors which produce the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof, and antibodies to these proteins and fragments thereof.



5 The present invention also provides methods for stimulating cellular proliferation by administering vectors, or cells containing vectors, encoded for production of the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The nucleotides and antibodies of the present invention are useful for the detection, localization and measurement of the nucleic acids encoding for the production of the proteins of the present invention, and also for the detection, localization and measurement of the proteins of the present invention. These nucleotides and antibodies can be combined with other reagents in kits for the purposes of detection, localization and measurement. These kits are useful for diagnosis and prognosis of conditions involving cellular proliferation associated with production of reactive oxygen intermediates.

10 The present invention solves the problems described above by providing a composition comprising the nucleotide sequence SEQ ID NO:1 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:3 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:22 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:41 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:45 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:47 and fragments thereof.

15 The present invention provides a composition comprising the protein SEQ ID NO:2 encoded by the nucleotide sequence SEQ ID NO:1. The present invention provides a composition comprising the protein SEQ ID NO:4 encoded by the nucleotide sequence SEQ ID NO:3. The present invention provides a composition comprising the protein SEQ ID NO:21 encoded by the nucleotide sequence SEQ ID NO:22. The present invention provides a composition comprising the protein



SEQ ID NO:42 encoded by the nucleotide sequence SEQ ID NO:41. The present invention provides a composition comprising the protein SEQ ID NO:46 encoded by the nucleotide sequence SEQ ID NO:45. The present invention provides a composition comprising the protein SEQ ID NO:48 encoded by the nucleotide sequence SEQ ID NO:47.

The present invention provides a composition comprising the protein SEQ ID NO:2 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:1 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:4 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:3 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:21 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:22 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:42 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:41 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:46 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:45 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:48 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:47 or fragments thereof.

The present invention also provides vectors containing the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. The present invention also provides cells transfected with these vectors. In addition, the present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:1 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof. The



present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:47 or fragments thereof.

The present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:1 or fragments thereof, which produce the protein SEQ ID NO:2 or fragments thereof. In addition, the present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof which produce the protein SEQ ID NO:4 or fragments thereof. In addition, the present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof which produce the protein SEQ ID NO:21 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof which produce the protein SEQ ID NO:42 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof which produce the protein SEQ ID NO:46 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:47 or fragments thereof which produce the protein SEQ ID NO:48 or fragments thereof.

The present invention provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:1 which produce the protein SEQ ID NO:2 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof, which produce the protein SEQ ID NO:4 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof, which produce



the protein SEQ ID NO:21 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof, which produce the protein SEQ ID NO:42 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof, which produce the protein SEQ ID NO:46 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:47 or fragments thereof, which produce the protein SEQ ID NO:48 or fragments thereof.

Specifically, the present invention provides a method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:1 or fragments thereof, which produce the protein SEQ ID NO:2 or fragments thereof. The present invention also provides a method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof, which produce the protein SEQ ID NO:4 or fragments thereof. The present invention also provides a method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof, which produce the protein SEQ ID NO:21 or fragments thereof. The present invention also provides a method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof, which produce the protein SEQ ID NO:42 or fragments thereof. The present invention also provides a method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof, which produce the protein SEQ ID NO:46 or fragments thereof. The present invention also provides a method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:47



or fragments thereof, which produce the protein SEQ ID NO:48 or fragments thereof.

The present invention may also be used to develop anti-sense nucleotide sequences to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. These anti-sense molecules may be used to interfere with translation of nucleotide sequences, such as SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins such as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof. Administration of these anti-sense molecules, or vectors encoding for anti sense molecules, to humans and animals, would interfere with production of proteins such as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, thereby decreasing production of ROIs and inhibiting cellular proliferation. These methods are useful in producing animal models for use in study of tumor development and vascular growth, and for study of the efficacy of treatments for affecting tumor and vascular growth *in vivo*.

The present invention also provides a method for high throughput screening of drugs and chemicals which modulate the proliferative activity of the enzymes of the present invention, thereby affecting cell division. Combinatorial chemical libraries may be screened for chemicals which modulate the proliferative activity of these enzymes. Drugs and chemicals may be evaluated based on their ability to modulate the enzymatic activity of the expressed or endogenous proteins, including those represented by SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof. Endogenous proteins may be obtained from many different tissues or cells, such as colon cells. Drugs may also be evaluated based on their ability to bind to the expressed or endogenous proteins represented by SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID



NO:46, SEQ ID NO:48 or fragments thereof. Enzymatic activity may be NADPH- or NADH-dependent superoxide generation catalyzed by the holoprotein. Enzymatic activity may also be NADPH- or NADH-dependent diaphorase activity catalyzed by either the holoprotein or the flavoprotein domain.

By flavoprotein domain, is meant approximately the C-terminal half of the enzymes shown in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, or fragments thereof, and the C-terminal end of the enzymes shown in SEQ ID NO:46 and SEQ ID NO:48 (approximately the C-terminal 265 amino acids). This fragment of gp91phox has NADPH-dependent reductase activity towards cytochrome c, nitrobluetetrazolium and other dyes. Expressed proteins or fragments thereof can be used for robotic screens of existing combinatorial chemical libraries. While not wanting to be bound by the following statement, it is believed that the NADPH or NADH binding site and the FAD binding site are useful for evaluating the ability of drugs and other compositions to bind to the mox and duox enzymes or to modulate their enzymatic activity. The use of the holoprotein or the C-terminal half or end regions are preferred for developing a high throughput drug screen. Additionally, the N-terminal one-third of the duox domain (the peroxidase domain) may also be used to evaluate the ability of drugs and other compositions to inhibit the peroxidase activity, and for further development of a high throughput drug screen.

The present invention also provides antibodies directed to the proteins SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The antibodies of the present invention are useful for a variety of purposes including localization, detection and measurement of the proteins SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The antibodies may be employed in kits to accomplish these purposes. These antibodies may also be linked to cytotoxic agents for selected killing of cells. The



term antibody is meant to include any class of antibody such as IgG, IgM and other classes. The term antibody also includes a completely intact antibody and also fragments thereof, including but not limited to Fab fragments and Fab + Fc fragments.

5           The present invention also provides the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 and fragments thereof. These nucleotides are useful for a variety of purposes including localization, detection, and measurement of messenger  
10       RNA involved in synthesis of the proteins represented as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. These nucleotides may also be used in the construction of labeled probes for the localization, detection, and measurement of  
15       nucleic acids such as messenger RNA or alternatively for the isolation of larger nucleotide sequences containing the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. These nucleotide sequences may  
20       be used to isolate homologous strands from other species using techniques known to one of ordinary skill in the art. These nucleotide sequences may also be used to make probes and complementary strands. In particular, the nucleotide sequence shown in SEQ ID NO:47 may be used to isolate the complete  
25       coding sequence for duox2. The nucleotides may be employed in kits to accomplish these purposes.

          Most particularly, the present invention involves a method for modulation of growth by modifying the proteins represented as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21,  
30       SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof.

          The term "mitogenic regulators" is used herein to mean any molecule that acts to affect cell division.

          The term "animal" is used herein to mean humans  
35       and non-human animals of both sexes.



The terms "a", "an" and "the" as used herein are defined to mean one or more and include the plural unless the context is inappropriate.

5 "Proteins", "peptides," "polypeptides" and "oligopeptides" are chains of amino acids (typically L-amino acids) whose alpha carbons are linked through peptide bonds formed by a condensation reaction between the carboxyl group of the alpha carbon of one amino acid and the amino group of the alpha carbon of another amino acid. The terminal amino acid at one end of the chain (*i.e.*, the amino terminal) has a free amino group, while the terminal amino acid at the other end of the chain (*i.e.*, the carboxy terminal) has a free carboxyl group. As such, the term "amino terminus" (abbreviated N-terminus) refers to the free alpha-amino group on the amino acid at the amino terminal of the protein, or to the alpha-amino group (imino group when participating in a peptide bond) of an amino acid at any other location within the protein. Similarly, the term "carboxy terminus" (abbreviated C-terminus) refers to the free carboxyl group on the amino acid at the carboxy terminus of a protein, or to the carboxyl group of an amino acid at any other location within the protein.

Typically, the amino acids making up a protein are numbered in order, starting at the amino terminal and increasing in the direction toward the carboxy terminal of the protein. Thus, when one amino acid is said to "follow" another, that amino acid is positioned closer to the carboxy terminal of the protein than the preceding amino acid.

The term "residue" is used herein to refer to an amino acid (D or L) or an amino acid mimetic that is incorporated into a protein by an amide bond. As such, the amino acid may be a naturally occurring amino acid or, unless otherwise limited, may encompass known analogs of natural amino acids that function in a manner similar to the naturally occurring amino acids (*i.e.*, amino acid mimetics). Moreover, an amide bond mimetic includes peptide backbone modifications well known to those skilled in the art.



Furthermore, one of skill will recognize that, as mentioned above, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 1%) in an encoded sequence are conservatively modified variations where the alterations result in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. The following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

When the peptides are relatively short in length (*i.e.*, less than about 50 amino acids), they are often synthesized using standard chemical peptide synthesis techniques. Solid phase synthesis in which the C-terminal amino acid of the sequence is attached to an insoluble support followed by sequential addition of the remaining amino acids in the sequence is a preferred method for the chemical synthesis of the antigenic epitopes described herein. Techniques for solid phase synthesis are known to those skilled in the art.

Alternatively, the antigenic epitopes described herein are synthesized using recombinant nucleic acid methodology. Generally, this involves creating a nucleic acid sequence that encodes the peptide or protein, placing the nucleic acid in an expression cassette under the control of a particular promoter, expressing the peptide or protein in a host, isolating the expressed peptide or protein and, if required, renaturing the



peptide or protein. Techniques sufficient to guide one of skill through such procedures are found in the literature.

When several desired protein fragments or peptides are encoded in the nucleotide sequence incorporated into a vector, one of skill in the art will appreciate that the protein fragments or peptides may be separated by a spacer molecule such as, for example, a peptide, consisting of one or more amino acids. Generally, the spacer will have no specific biological activity other than to join the desired protein fragments or peptides together, or to preserve some minimum distance or other spatial relationship between them. However, the constituent amino acids of the spacer may be selected to influence some property of the molecule such as the folding, net charge, or hydrophobicity. Nucleotide sequences encoding for the production of residues which may be useful in purification of the expressed recombinant protein may be built into the vector. Such sequences are known in the art. For example, a nucleotide sequence encoding for a poly histidine sequence may be added to a vector to facilitate purification of the expressed recombinant protein on a nickel column.

Once expressed, recombinant peptides, polypeptides and proteins can be purified according to standard procedures known to one of ordinary skill in the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like. Substantially pure compositions of about 50 to 99% homogeneity are preferred, and 80 to 95% or greater homogeneity are most preferred for use as therapeutic agents.

One of skill in the art will recognize that after chemical synthesis, biological expression or purification, the desired proteins, fragments thereof and peptides may possess a conformation substantially different than the native conformations of the proteins, fragments thereof and peptides. In this case, it is often necessary to denature and reduce protein and then to cause the protein to re-fold into the preferred conformation. Methods of reducing and denaturing proteins



and inducing re-folding are well known to those of skill in the art.

5           The genetic constructs of the present invention include coding sequences for different proteins, fragments thereof, and peptides. The genetic constructs also include epitopes or domains chosen to permit purification or detection of the expressed protein. Such epitopes or domains include DNA sequences encoding the glutathione binding domain from glutathione S-transferase, hexa-histidine, thioredoxin, 10 hemagglutinin antigen, maltose binding protein, and others commonly known to one of skill in the art. The preferred genetic construct includes the nucleotide sequences of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. It is to be understood that additional or alternative nucleotide sequences may be included in the genetic constructs in order to encode for the following: a) multiple copies of the desired proteins, fragments thereof, or peptides; b) various combinations of the desired proteins, fragments thereof, or peptides; and c) 15 conservative modifications of the desired proteins, fragments thereof, or peptides, and combinations thereof. Preferred proteins include the human mox1 protein and human mox2 protein shown as SEQ ID NO:2 and SEQ ID NO:4, respectively, and fragments thereof. Some preferred fragments of the human mox1 protein (SEQ ID NO:2) include but are not limited to the proteins shown as SEQ ID NO:23, SEQ ID NO:24, and SEQ ID NO:25. The protein mox1 is also called p65mox in this application. Another preferred protein of the present invention is rat mox1 protein shown as SEQ ID NO:21 and fragments thereof. Another preferred protein of the present invention is rat mox1B protein shown as SEQ ID NO:42 and fragments thereof. Yet another preferred protein of the present invention is duox1 protein shown as SEQ ID NO:46 and fragments thereof. Still another preferred protein of the present invention is duox2 protein. A partial amino acid sequence of the duox2 protein is shown as SEQ ID NO:48. 20 25 30 35



The nucleotide sequences of the present invention may also be employed to hybridize to nucleic acids such as DNA or RNA nucleotide sequences under high stringency conditions which permit detection, for example, of alternately spliced messages.

The genetic construct is expressed in an expression system such as in NIH 3T3 cells using recombinant sequences in a pcDNA-3 vector (Invitrogen, Carlsbad, CA) to produce a recombinant protein. Preferred expression systems include but are not limited to Cos-7 cells, insect cells using recombinant baculovirus, and yeast. It is to be understood that other expression systems known to one of skill in the art may be used for expression of the genetic constructs of the present invention. The preferred proteins of the present invention are the proteins referred to herein as human mox1 and human mox2 or fragments thereof which have the amino acid sequences set forth in SEQ ID NO:3 and SEQ ID NO:4, respectively, or an amino acid sequence having amino acid substitutions as defined in the definitions that do not significantly alter the function of the recombinant protein in an adverse manner. Another preferred protein of the present invention is referred to herein as rat mox1 and has the amino acid sequence set forth in SEQ ID NO:21. Yet another preferred protein of the present invention is referred to herein as rat mox1B and has the amino acid sequence set forth in SEQ ID NO:42. Two other preferred proteins of the present invention are referred to herein as human duox1 and human duox2, or fragments thereof, which have the amino acid sequences set forth in SEQ ID NO:46 and SEQ ID NO:48, respectively, or an amino acid sequence having amino acid substitutions as defined in the definitions that do not significantly alter the function of the recombinant protein in an adverse manner.

#### *Terminology*

It should be understood that some of the terminology used to describe the novel mox and duox proteins



contained herein is different from the terminology in U.S. Provisional Application Serial No. 60/107,911 and U.S. Provisional Application Serial No. 60/149,332 upon which this application claims priority in part. As described herein, the term "human mox1" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:2, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:1, or a fragment thereof. As described herein, the term "human mox2" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:4, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:3, or a fragment thereof. As described herein, the term "human duox1" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:46, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:45, or a fragment thereof. As described herein, the term "human duox2" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:48, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:47, or a fragment thereof.

#### *Construction of the Recombinant Gene*

The desired gene is ligated into a transfer vector, such as pcDNA3, and the recombinants are used to transform host cells such as Cos-7 cells. It is to be understood that different transfer vectors, host cells, and transfection methods may be employed as commonly known to one of ordinary skill in the art. Six desired genes for use in transfection are shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45 and SEQ ID NO:47. For example, lipofectamine-mediated transfection and *in vivo* homologous recombination was used to introduce the mox1 gene into NIH 3T3 cells.

The synthetic gene is cloned and the recombinant construct containing mox or duox gene is produced and grown in confluent monolayer cultures of a Cos-7 cell line. The



expressed recombinant protein is then purified, preferably using affinity chromatography techniques, and its purity and specificity determined by known methods.

5 A variety of expression systems may be employed for expression of the recombinant protein. Such expression methods include, but are not limited to the following: bacterial expression systems, including those utilizing *E. coli* and *Bacillus subtilis*; virus systems; yeast expression systems; cultured insect and mammalian cells; and other expression systems known to  
10 one of ordinary skill in the art.

#### *Transfection of Cells*

It is to be understood that the vectors of the present invention may be transfected into any desired cell or cell line.  
15 Both *in vivo* and *in vitro* transfection of cells are contemplated as part of the present invention. Preferred cells for transfection include but are not limited to the following: fibroblasts (possibly to enhance wound healing and skin formation), granulocytes (possible benefit to increase function in a  
20 compromised immune system as seen in AIDS, and aplastic anemia), muscle cells, neuroblasts, stem cells, bone marrow cells, osteoblasts, B lymphocytes, and T lymphocytes.

Cells may be transfected with a variety of methods known to one of ordinary skill in the art and include but are not  
25 limited to the following: electroporation, gene gun, calcium phosphate, lipofectamine, and fugene, as well as adenoviral transfection systems.

Host cells transfected with the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22,  
30 SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, or fragments thereof, are used to express the proteins SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 and SEQ ID NO:48, respectively, or fragments thereof.

These expressed proteins are used to raise  
35 antibodies. These antibodies may be used for a variety of applications including but not limited to immunotherapy against



cancers expressing one of the mox or duox proteins, and for detection, localization and measurement of the proteins shown in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48 or fragments thereof.

5

#### *Purification and Characterization of the Expressed Protein*

The proteins of the present invention can be expressed as a fusion protein with a poly histidine component, such as a hexa histidine, and purified by binding to a metal affinity column using nickel or cobalt affinity matrices. The protein can also be expressed as a fusion protein with glutathione S-transferase and purified by affinity chromatography using a glutathione agarose matrix. The protein can also be purified by immunoaffinity chromatography by expressing it as a fusion protein, for example with hemagglutinin antigen. The expressed or naturally occurring protein can also be purified by conventional chromatographic and purification methods which include anion and cation exchange chromatography, gel exclusion chromatography, hydroxylapatite chromatography, dye binding chromatography, ammonium sulfate precipitation, precipitation in organic solvents or other techniques commonly known to one of skill in the art.

10

15

20

25

#### *Methods of Assessing Activity of Expressed Proteins*

Different methods are available for assessing the activity of the expressed proteins of the present invention, including but not limited to the proteins represented as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48 substituted analogs thereof, and fragments thereof.

30

#### 1. Assays of the holoprotein and fragments thereof for superoxide generation:

35

##### A. General considerations.



These assays are useful in assessing efficacy of drugs designed to modulate the activity of the enzymes of the present invention. The holoprotein may be expressed in COS-7 cells, NIH 3T3 cells, insect cells (using baculoviral technology) or other cells using methods known to one of skill in the art. Membrane fractions or purified protein are used for the assay. The assay may require or be augmented by other cellular proteins such as p47phox, p67phox, and Rac1, as well as potentially other unidentified factors (e.g., kinases or other regulatory proteins).

#### B. Cytochrome c reduction.

NADPH or NADH is used as the reducing substrate, in a concentration of about 100  $\mu$ M. Reduction of cytochrome c is monitored spectrophotometrically by the increase in absorbance at 550 nm, assuming an extinction coefficient of 21  $\text{mM}^{-1}\text{cm}^{-1}$ . The assay is performed in the absence and presence of about 10  $\mu$ g superoxide dismutase. The superoxide-dependent reduction is defined as cytochrome c reduction in the absence of superoxide dismutase minus that in the presence of superoxide dismutase (Uhlinger et al. (1991) *J. Biol. Chem.* 266, 20990-20997). Acetylated cytochrome c may also be used, since the reduction of acetylated cytochrome c is thought to be exclusively via superoxide.

#### C. Nitroblue tetrazolium reduction.

For nitroblue tetrazolium (NBT) reduction, the same general protocol is used, except that NBT is used in place of cytochrome c. In general, about 1 mL of filtered 0.25 % nitrotetrazolium blue (Sigma, St. Louis, MO) is added in Hanks buffer without or with about 600 Units of superoxide dismutase (Sigma) and samples are incubated at approximately 37°C. The oxidized NBT is clear, while the reduced NBT is blue and insoluble. The insoluble product is collected by centrifugation, and the pellet is re-suspended in about 1 mL of pyridine (Sigma) and heated for about 10 minutes at 100°C to solubilize



the reduced NBT. The concentration of reduced NBT is determined by measuring the absorbance at 510 nm, using an extinction coefficient of  $11,000 \text{ M}^{-1}\text{cm}^{-1}$ . Untreated wells are used to determine cell number.

5

#### D. Luminescence.

Superoxide generation may also be monitored with a chemiluminescence detection system utilizing lucigenin (bis-N-methylacridinium nitrate, Sigma, St. Louis, MO). The sample is mixed with about  $100 \mu\text{M}$  NADPH (Sigma, St. Louis, MO) and  $10 \mu\text{M}$  lucigenin (Sigma, St. Louis, MO) in a volume of about  $150 \mu\text{L}$  Hanks solution. Luminescence is monitored in a 96-well plate using a LumiCounter (Packard, Downers Grove, IL) for 0.5 second per reading at approximately 1 minute intervals for a total of about 5 minutes; the highest stable value in each data set is used for comparisons. As above, superoxide dismutase is added to some samples to prove that the luminescence arises from superoxide. A buffer blank is subtracted from each reading (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321).

20

#### E. Assays in intact cells.

Assays for superoxide generation may be performed using intact cells, for example, the mox-transfected NIH 3T3 cells. In principle, any of the above assays can be used to evaluate superoxide generation using intact cells, for example, the mox-transfected NIH 3T3 cells. NBT reduction is a preferred assay method.

25

30

#### 2. Assays of truncated proteins comprised of approximately the C-terminal 265 amino acid residues

35

While not wanting to be bound by the following statement, the truncated protein comprised of approximately the C-terminal 265 amino acid residues is not expected to generate superoxide, and therefore, superoxide dismutase is not added in assays of the truncated protein. Basically, a similar assay is



established and the superoxide-independent reduction of NBT, cytochrome c, dichlorophenolindophenol, ferricyanide, or another redox-active dye is examined.

5        *Nucleotides and Nucleic Acid Probes*

          The nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof and PCR primers therefor, may be used, respectively, for localization, detection  
10        and measurement of nucleic acids related to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof. The nucleotide sequences SEQ ID NO:1 and SEQ ID NO:3 are also called the human mox1 gene and the human mox2 gene in this application.  
15        SEQ ID NO:22 is also known as the rat mox1 gene in this application. SEQ ID NO:41 is also known as the rat mox1B gene in this application. SEQ ID NO:45 is also known as the human duox1 gene in this application. SEQ ID NO:47 is also known as the human duox2 gene in this application.

20        The nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof, may be used to create probes to isolate larger nucleotide sequences containing the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID  
25        NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, respectively. The nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof, may also be used to create probes to identify and isolate mox and duox proteins in  
30        other species.

          The nucleic acids described herein include messenger RNA coding for production of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. Such nucleic acids  
35        include but are not limited to cDNA probes. These probes may be labeled in a variety of ways known to one of ordinary skill in



the art. Such methods include but are not limited to isotopic and non-isotopic labeling. These probes may be used for *in situ* hybridization for localization of nucleic acids such as mRNA encoding for SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. Localization may be performed using *in situ* hybridization at both ultrastructural and light microscopic levels of resolution using techniques known to one of ordinary skill in the art.

These probes may also be employed to detect and quantitate nucleic acids and mRNA levels using techniques known to one of ordinary skill in the art including but not limited to solution hybridization.

#### *Antibody Production*

The proteins shown in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, are combined with a pharmaceutically acceptable carrier or vehicle to produce a pharmaceutical composition and administered to animals for the production of polyclonal antibodies using methods known to one of ordinary skill in the art. The preferred animals for antibody production are rabbits and mice. Other animals may be employed for immunization with these proteins or fragments thereof. Such animals include, but are not limited to the following; sheep, horses, pigs, donkeys, cows, monkeys and rodents such as guinea pigs and rats.

The terms "pharmaceutically acceptable carrier or pharmaceutically acceptable vehicle" are used herein to mean any liquid including but not limited to water or saline, oil, gel, salve, solvent, diluent, fluid ointment base, liposome, micelle, giant micelle, and the like, which is suitable for use in contact with living animal or human tissue without causing adverse physiological responses, and which does not interact with the other components of the composition in a deleterious manner.



The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets commonly used by one of ordinary skill in the art.

Preferred unit dosage formulations are those containing a dose or unit, or an appropriate fraction thereof, of the administered ingredient. It should be understood that in addition to the ingredients, particularly mentioned above, the formulations of the present invention may include other agents commonly used by one of ordinary skill in the art.

The pharmaceutical composition may be administered through different routes, such as oral, including buccal and sublingual, rectal, parenteral, aerosol, nasal, intramuscular, subcutaneous, intradermal, and topical. The pharmaceutical composition of the present invention may be administered in different forms, including but not limited to solutions, emulsions and suspensions, microspheres, particles, microparticles, nanoparticles, and liposomes. It is expected that



from about 1 to 7 dosages may be required per immunization regimen. Initial injections may range from about 0.1  $\mu$ g to 1 mg, with a preferred range of about 1  $\mu$ g to 800  $\mu$ g, and a more preferred range of from approximately 25  $\mu$ g to 500  $\mu$ g. Booster injections may range from 0.1  $\mu$ g to 1 mg, with a preferred range of approximately 1  $\mu$ g to 800  $\mu$ g, and a more preferred range of about 10  $\mu$ g to 500  $\mu$ g.

The volume of administration will vary depending on the route of administration and the size of the recipient. For example, intramuscular injections may range from about 0.1 ml to 1.0 ml.

The pharmaceutical composition may be stored at temperatures of from about 4°C to -100°C. The pharmaceutical composition may also be stored in a lyophilized state at different temperatures including room temperature. The pharmaceutical composition may be sterilized through conventional means known to one of ordinary skill in the art. Such means include, but are not limited to filtration, radiation and heat. The pharmaceutical composition of the present invention may also be combined with bacteriostatic agents, such as thimerosal, to inhibit bacterial growth.

#### *Adjuvants*

A variety of adjuvants known to one of ordinary skill in the art may be administered in conjunction with the protein in the pharmaceutical composition. Such adjuvants include, but are not limited to the following: polymers, copolymers such as polyoxyethylene-polyoxypropylene copolymers, including block co-polymers; polymer P1005; Freund's complete adjuvant (for animals); Freund's incomplete adjuvant; sorbitan monooleate; squalene; CRL-8300 adjuvant; alum; QS 21, muramyl dipeptide; trehalose; bacterial extracts, including mycobacterial extracts; detoxified endotoxins; membrane lipids; or combinations thereof.

Monoclonal antibodies can be produced using hybridoma technology in accordance with methods well known



to those skilled in the art. The antibodies are useful as research or diagnostic reagents or can be used for passive immunization. The composition may optionally contain an adjuvant.

5 The polyclonal and monoclonal antibodies useful as research or diagnostic reagents may be employed for detection and measurement of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. Such antibodies may be used to detect these  
10 proteins in a biological sample, including but not limited to samples such as cells, cellular extracts, tissues, tissue extracts, biopsies, tumors, and biological fluids. Such detection capability is useful for detection of disease related to these proteins to facilitate diagnosis and prognosis and to suggest possible treatment alternatives.

15 Detection may be achieved through the use of immunocytochemistry, ELISA, radioimmunoassay or other assays as commonly known to one of ordinary skill in the art. The mox1, mox2, duox1 and duox2 proteins, or fragments thereof, may be labeled through commonly known approaches,  
20 including but not limited to the following: radiolabeling, dyes, magnetic particles, biotin-avidin, fluorescent molecules, chemiluminescent molecules and systems, ferritin, colloidal gold, and other methods known to one of skill in the art of labeling proteins.

#### 25 *Administration of Antibodies*

The antibodies directed to the proteins shown as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48, or directed to fragments  
30 thereof, may also be administered directly to humans and animals in a passive immunization paradigm. Antibodies directed to extracellular portions of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48 bind to these extracellular epitopes. Attachment of  
35 labels to these antibodies facilitates localization and visualization of sites of binding. Attachment of molecules such as ricin or



other cytotoxins to these antibodies helps to selectively damage or kill cells expressing SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof.

5

### *Kits*

The present invention includes kits useful with the antibodies, nucleic acids, nucleic acid probes, labeled antibodies, labeled proteins or fragments thereof for detection, localization and measurement of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48 or combinations and fragments thereof.

10

Kits may be used for immunocytochemistry, *in situ* hybridization, solution hybridization, radioimmunoassay, ELISA, Western blots, quantitative PCR, and other assays for the detection, localization and measurement of these nucleic acids, proteins or fragments thereof using techniques known to one of skill in the art.

15

The nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, may also be used under high stringency conditions to detect alternately spliced messages related to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45, SEQ ID NO:47 or fragments thereof, respectively.

20

As discussed in one of the Examples, rat mox1 protein (SEQ ID NO: 21) is similar to mouse gp91 protein (SEQ ID NO: 38), whereas rat mox1B protein (SEQ ID NO:42) is similar to human gp91 protein (SEQ ID NO:12). This observation suggests that other isoforms of mouse and human gp91 may exist. In addition, another subtype of human mox1, similar to rat mox1B (SEQ ID NO:42), also exists. The presence of two isoforms of rat mox1 protein in vascular smooth muscle may have important physiological consequences

25

30

35



and biomedical applications. For example, the two isoforms may have different biological activities, different tissue distributions and may be regulated differently in physiological and/or pathological conditions. The fact that mox1B (SEQ ID NO:42) was isolated from cells exposed to angiotensin II, known to promote oxidative stress and vascular growth, suggests that it may be upregulated by this hormone and may be overexpressed in disease. Therefore, the diagnostic kits of the present invention can measure the relative expression of the two mox isoforms. The diagnostic kits may also measure or detect the relative expression of the mox proteins described herein (i.e. human mox1 and/or human mox2) and duox proteins described herein (i.e. human duox1 and/or human duox2).

Fragments of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47 containing the relevant hybridizing sequence can be synthesized onto the surface of a chip array. RNA samples, e.g., from tumors, are then fluorescently tagged and hybridized onto the chip for detection. This approach may be used diagnostically to characterize tumor types and to tailor treatments and/or provide prognostic information. Such prognostic information may have predictive value concerning disease progression and life span, and may also affect choice of therapy.

The present invention is further illustrated by the following examples, which are not to be construed in any way as imposing limitations upon the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other embodiments, modifications, and equivalents thereof, which, after reading the description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention.



## EXAMPLE 1

*Sequence Analysis and Cloning of the Human mox1 cDNA (SEQ ID NO:1) Encoding for Production of the Human mox1 Protein p65mox (SEQ ID NO:2)*

5           Using gp91phox as a query sequence, a 334 base sequenced portion of expressed sequence tag (EST) 176696 (GenBank Accession number AA305700) showed 68.8% sequence identity at the predicted amino acid level with human (h)-gp91phox. The bacterial strain number 129134 containing the EST sequence in the pBluescript SK<sup>-</sup> vector, was purchased from American Tissue Type Culture Collection (ATCC, Rockville, MD). The EST sequence was originally cloned from a Caco-2 human colon carcinoma cell line. The EST176696 DNA was further sequenced using the T7 and T3 vector promoters and primers designed to match the known 3' sequence. Internal primers used for sequencing were as follows: 5'-AAC AAG CGT GGC TTC AGC ATG-3' SEQ ID NO:5 (251S, numbering is based on the nucleotides from the 5' end of EST176696, and S indicates the sense direction), 5'-AGC AAT ATT GTT GGT CAT-3' SEQ ID NO:6 (336S), 5'-GAC TTG ACA GAA AAT CTA TAA GGG-3' SEQ ID NO:7 (393S), 5'-TTG TAC CAG ATG GAT TTC AA-3' SEQ ID NO:8 (673A, A indicates the antisense direction), 5'-CAG GTC TGA AAC AGA AAA CCT-3' SEQ ID NO:9 (829S), 5'-ATG AAT TCT CAT TAA TTA TTC AAT AAA-3' SEQ ID NO:10 (1455A). The coding sequence in EST176696 showed homology to a 250 amino acid stretch corresponding to the N-terminal 44% of human gp91phox, and contained a stop codon corresponding to the location in human gp91phox. 5' Rapid amplification of cDNA ends (RACE) was carried out using a human colon cDNA library and Marathon cDNA Amplification Kit (ClonTech) using 5'-ATC TCA AAA GAC TCT GCA CA-3' SEQ ID NO:11 (41A) as an internal gene-specific primer (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85, 8998-9002). 5' RACE resulted in a 1.1 kb fragment representing the complete 5' sequence, based on homology with gp91phox.



Reamplification was performed with primers spanning the putative start and stop codons, using the 1.1 kb 5' RACE product and pSK-EST176696 for primer design. The amplified 1.7kb fragment was TA cloned into the PCR2.1 vector (Invitrogen, Carlsbad, CA). This recombinant vector is referred to as PCR-*mox*.

Figure 1(a-d) presents a comparison of the present amino acid sequences of human, bovine and murine gp91 phox with the human and rat *mox1* proteins of the present invention and the human *duox2* protein of the present invention. Also shown are the amino acid sequences for related plant enzyme proteins.

The encoded hp65mox ("mox" referring to mitogenic oxidase and "65" referring to its predicted molecular weight) is listed as SEQ ID NO:2. h-gp91phox (SEQ ID NO:12) and SEQ ID NO:2 differ in length by 3 residues and are 70% identical in their amino acid sequence. h-gp91phox and SEQ ID NO:2 show a greater percentage identity in the C-terminal half of the molecule which contains the putative NADPH and FAD binding sites, and there are several relatively long stretches of complete identity within this region.

A dendrogram (Figure 2) comparing the amino acid sequences of mouse and human gp91phox with that of *mox1* SEQ ID NO:2 shows that the latter probably represents a distinct isoform of gp91phox. Two plant homologs of cytochrome  $b_{558}$  large subunit are also indicated and represent more distant relatives of the human sequences. Human (and rat *mox1* described more fully below) lack asparagine-linked glycosylation sites, which are seen in the highly glycosylated human and mouse gp91phox. Additionally, the hydropathy profiles of human gp91phox and *mox1* are nearly identical and include five very hydrophobic stretches in the amino-terminal half of the molecules which are predicted to be membrane-spanning regions.



## EXAMPLE 2

### *Expression of Mox1*

Human multiple tissue northern (MTN) Blot I and Human MTN Blot IV (ClonTech) membranes were hybridized with the putative coding region of the PCR-mox vector at 68°C for several hours. The mox coding region was labeled by random priming with [ $\alpha$ -<sup>32</sup>P]dCTP (10  $\mu$ Ci) using the Prime-It II kit (Stratagene). For analysis of mox1 expression in cell lines, total RNA was prepared from 10<sup>6</sup> cells using the High Pure RNA Isolation Kit (Boehringer Mannheim) or RNeasy kit (Quiagen). Total RNA (10-20  $\mu$ g) was separated on a 1% agarose formaldehyde mini-gel and transferred to a Nytran filter (Biorad) and immobilized by ultraviolet cross-linking.

Northern blotting revealed that the major location of mRNA coding for the mox1 protein was colon. The message was also detected in prostate and uterus. The human colon-carcinoma cell line, Caco-2, also expressed large quantities of mox1 message. Northern blotting of mRNA from rat aortic smooth muscle cells also showed strong hybridization, which increased roughly two-fold within 12 hours after treatment with platelet-derived growth factor. This increase in the expression of rat mox1 is consistent with the idea that mox1 contributes to the growth-stimulatory effects of PDGF.

## EXAMPLE 3

### *Transfection of NIH3T3 Cells with SEQ ID NO:1*

The nucleotide sequence (SEQ ID NO:1) encoding for production of the mox1 protein (SEQ ID NO:2) was subcloned into the *NotI* site of the pEF-PAC vector (obtained from Mary Dinuer, Indiana University Medical School, Indianapolis, IN) which has a puromycin resistance gene. Transfection was carried out as described in Sambrook et al., Molecular Cloning, A Laboratory Manual, Volumes 1-3, 2nd edition, Cold Spring Harbor Laboratory Press, N.Y., 1989. The SEQ ID NO:1 in pEF-PAC and the empty vector were separately transfected into NIH 3T3 cells using Fugene 6



(Boeringer Mannheim). About  $2 \times 10^6$  cells maintained in DMEM containing 10% calf serum were transfected with 10  $\mu$ g of DNA. After 2 days, cells were split and selected in the same medium containing 1mg/ml puromycin. Colonies that survived in selection media for 10 to 14 days were subcultured continuously in the presence of puromycin.

Transfected cells exhibited a "transformed"-like morphology, similar to that seen with (V12)Ras-transfected cells, characterized by long spindle-like cells. The parent NIH 3T3 cells or cells transfected with the empty vector showed a normal fibroblast-like morphology.

#### EXAMPLE 4

*Expression of Mox1 (SEQ ID NO:1) in Transfected NIH3T3 Cells*

To verify the expression of *mox1* mRNA after transfection, RT-PCR and Northern blotting were performed. Total RNAs were prepared from  $10^6$  cells using the High Pure RNA Isolation Kit (Boeringer Mannheim) or RNeasy kit (Qiagen). cDNAs for each colony were prepared from 1-2  $\mu$ g of total RNA using Advantage RT-PCR Kit (ClonTech). PCR amplification was performed using primers, 5'-TTG GCT AAA TCC CAT CCA-3' SEQ ID NO:13 (NN459S, numbering containing NN indicates numbering from the start codon of *mox1*) and 5'-TGC ATG ACC AAC AAT ATT GCT G-3' SEQ ID NO:14 (NN1435A). For Northern blotting, 10-20  $\mu$ g of total RNA was separated on a 1% agarose formaldehyde gel and transferred to a nylon filter. After ultraviolet (UV) cross-linking, filters were used for Northern blotting assay as described in Example 2.

Colonies expressing large amounts of *mox1* mRNA were chosen for further analysis. The expression of mRNA for glyceraldehyde 3 phosphate dehydrogenase in the various cell lines was normal.

#### EXAMPLE 5



### *Colony Formation on Soft Agar*

10<sup>5</sup> to 10<sup>3</sup> cells stably transfected with human  
mox1 gene SEQ ID NO:1 and with empty vector were prepared  
in 0.3% warm (40°C) agar solution containing DMEM and 10%  
5 calf serum. Cells were distributed onto a hardened 0.6% agar  
plate prepared with DMEM and 10% calf serum. After three  
weeks in culture (37°C, 5% CO<sub>2</sub>) colony formation was  
observed by microscopy.

10 Cells which were stably transfected with the empty  
vector and cultured in soft agar for 3 weeks as above did not  
display anchorage independent growth. In contrast, NIH 3T3  
cells which had been stably transfected with the mox1 (SEQ ID  
NO:1) and cultured for 3 weeks in soft agar demonstrated  
anchorage independent growth of colonies.

### **EXAMPLE 6**

#### *NADPH-Dependent Superoxide Generation Assay*

In one embodiment of the present invention, NIH  
3T3 cells stably transfected with the human mox1 gene (SEQ ID  
20 NO:1) were analyzed for superoxide generation using the  
lucigenin (Bis-N-methylacridinium luminescence assay (Sigma,  
St. Louis, MO, Li et al. (1998) *J. Biol. Chem.* 273, 2015-  
2023). Cells were washed with cold HANKS' solution and  
homogenized on ice in HANKS' buffer containing 15% sucrose  
25 using a Dounce homogenizer. Cell lysates were frozen  
immediately in a dry ice/ethanol bath. For the assay, 30 µg of  
cell lysate was mixed with 200 µM NADPH and 500 µM  
lucigenin. Luminescence was monitored using a LumiCounter  
(Packard) at three successive one minute intervals and the  
30 highest value was used for comparison. Protein concentration  
was determined by the Bradford method.

Superoxide generation was monitored in lysates  
from some of the stably transfected cell lines and was compared  
with superoxide generation by the untransfected NIH 3T3 cell  
35 lysates. The results are shown in Table 4. Cell lines 26, 27,  
and 28 gave the highest degree of morphological changes by



microscopic examination corresponding to the highest degree of superoxide generation. The luminescent signal was inhibited by superoxide dismutase and the general flavoprotein inhibitor diphenylene iodonium, but was unaffected by added recombinant human p47phox, p67phox and Rac1(GTP- $\gamma$ S), which are essential cytosolic factors for the phagocyte respiratory-burst oxidase.

**Table 4**

10	Cell Line Name	Superoxide Generation (RLU)
	Control (untransfected)	6045
	mox1-26	17027
	mox1-27	14670
15	mox1-28	18411
	mox1-65	5431
	mox1-615	11331
	mox1-+3	8645
	mox1-+10	5425
20	mox1-pcc16	8050

In an alternate and preferred embodiment of the present invention, cells that had been stably transfected with mox1 (YA28) or with empty vector (NEF2) were grown in 10 cm tissue culture plates in medium containing DMEM, 10% calf serum, 100 units/ml penicillin, 100  $\mu$ g/ml streptomycin, and 1  $\mu$ g/ml puromycin to approximately 80% confluency. Cells (five tissue culture plates of each cell type) were washed briefly with 5 ml phosphate buffered saline (PBS) then dissociated from the plates with PBS containing 5 mM EDTA. Cells were pelleted by centrifuging briefly at 1000 x g.

To permeabilize the cells, freeze thaw lysis was carried out and this was followed by passage of the cell material through a small bore needle. The supernatant was removed and the cells were frozen on dry ice for 15 minutes. After cells were thawed, 200  $\mu$ l lysis buffer (HANKS' Buffered Salt



Solution - HBBS) containing a mixture of protease inhibitors from Sigma (Catalog # P2714) was added. Cells on ice were passed through an 18 gauge needle 10 times and 200  $\mu$ l of HBSS buffer containing 34% sucrose was added to yield a final concentration of 17% sucrose. Sucrose appeared to enhance stability upon storage. The combination of freeze-thawing and passage through a needle results in lysis of essentially all of the cells, and this material is referred to as the "cell lysate."

The cell lysates were assayed for protein concentration using the BioRad protein assay system. Cell lysates were assayed for NADPH-dependent chemiluminescence by combining HBSS buffer, arachidonic acid, and 0.01 - 1  $\mu$ g protein in assay plates (96 well plastic plates). The reaction was initiated by adding 1.5 mM NADPH and 75  $\mu$ M lucigenin to the assay mix to give a final concentration of 200  $\mu$ M NADPH and 10  $\mu$ M lucigenin, and the chemiluminescence was monitored immediately. The final assay volume as 150  $\mu$ l. The optimal arachidonic acid concentration was between 50-100  $\mu$ M. A Packard Lumicount luminometer was used to measure chemiluminescence of the reaction between lucigenin and superoxide at 37°C. The plate was monitored continuously for 60 minutes and the maximal relative luminescence unit (RLU) value for each sample was used for the graph.

Figure 3 shows the RLU at various concentrations of cell lysates from mox1-transfected (YA28) and vector control (NEF2) cells. The presence of NaCl or KCl within a concentration range of 50-150  $\mu$ M is important for optimal activity.  $MgCl_2$  (1-5 mM) further enhanced activity by about 2-fold. This cell-free assay for mox1 NADPH-oxidase activity is useful for screening modulators (inhibitors or stimulators) of the mox1 enzyme. The assay may also be used to detect mox and duox NADPH-oxidase activity in general and to screen for modulators (inhibitors or stimulators) of the mox and duox family of enzymes.



**EXAMPLE 7**

*Nitro Blue Tetrazolium Reduction by Superoxide Generated by NIH 3T3 cells Transfected with the Mox1 cDNA (SEQ ID NO:1)*

5                   Superoxide generation by intact cells was monitored by using superoxide dismutase-sensitive reduction of nitroblue tetrazolium. NEF2 (vector alone control), YA26 (mox1 (SEQ ID NO:1)-transfected) and YA28 (mox1 (SEQ ID NO:1)-transfected) cells were plated in six well plates at 10                   500,000 cells per well. About 24 hours later, medium was removed from cells and the cells were washed once with 1 mL Hanks solution (Sigma, St. Louis, MO). About 1 mL of filtered 0.25% Nitro blue tetrazolium (NBT, Sigma) was added in Hanks without or with 600 units of superoxide dismutase 15                   (Sigma) and cells were incubated at 37°C in the presence of 5% CO<sub>2</sub>. After 8 minutes the cells were scraped and pelleted at more than 10,000g. The pellet was re-suspended in 1 mL of pyridine (Sigma) and heated for 10 minutes at 100°C to solubilize the reduced NBT. The concentration of reduced NBT 20                   was determined by measuring the absorbance at 510 nm, using an extinction coefficient of 11,000 M<sup>-1</sup>cm<sup>-1</sup>. Some wells were untreated and used to determine cell number.

25                   The data are presented in Table 5 and Figure 4 and indicate that the mox1 (SEQ ID NO:1)-transfected cells generated significant quantities of superoxide.

**Table 5**

	<u>NBT Reduction (nmols/10<sup>6</sup> cells)</u>	<u>- SOD</u>	<u>+ SOD</u>
30	vector control cells	2.5 ± 0.5	2.1 ± 0.5
	YA26 (mox1) cells	6.4 ± 0.2	3.4 ± 0.1
	YA28 (mox1) cells	5.2 ± 0.6	3.4 ± 0.3

35                   -SOD, and +SOD mean in the absence or presence of added superoxide dismutase, respectively.

Because superoxide dismutase is not likely to penetrate cells, superoxide must be generated extracellularly. The amount of



superoxide generated by these cells is about 5-10% of that generated by activated human neutrophils.

#### EXAMPLE 8

##### 5      *Modification of Intracellular Components in Mox1 Transfected Cells*

To test whether superoxide generated by *mox1* can affect intracellular "targets," aconitase activity in control and *mox*-transfected cell lines was monitored as described in Suh et al.  
10      (1999) *Nature* 401, 79-82. Aconitase contains a four-iron-sulphur cluster that is highly susceptible to modification by superoxide, resulting in a loss of activity, and has been used as a reporter of intra-cellular superoxide generation. Aconitase activity was determined as described in Gardner et al. (1995) *J.*  
15      *Biol. Chem.* 270, 13399-13405. Aconitase activity was significantly diminished in all three *mox*-transfected cell lines designated YA26, YA28 and YA212 as compared to the transfected control (Figure 5). Approximately 50% of the aconitase in these cells is mitochondrial, based on differential  
20      centrifugation, and the cytosolic and mitochondrial forms were both affected. Control cytosolic and mitochondrial enzymes that do not contain iron-sulfur centres were not affected. Superoxide generated in *mox1*-transfected cells is therefore capable of reacting with and modifying intracellular  
25      components.

#### EXAMPLE 9

##### *Tumor Generation in Nude Mice Receiving Cells Transfected with the Human *mox1* cDNA (SEQ ID NO:1)*

30      About  $2 \times 10^6$  NIH 3T3 cells (either *mox1*-transfected with SEQ ID NO:1 or cells transfected using empty vector) were injected subdermally into the lateral aspect of the neck of 4-5 week old nude mice. Three to six mice were injected for each of three *mox1*-transfected cell lines, and 3  
35      mice were injected with the cells transfected with empty vector (control). After 2 to 3 weeks, mice were sacrificed. The tumors were fixed in 10% formalin and characterized by



histological analysis. Tumors averaged 1.5 x 1 x 1 cm in size and showed histology typical of sarcoma type tumors. In addition, tumors appeared to be highly vascularized with superficial capillaries. Eleven of twelve mice injected with  
5 mox1 gene-transfected cells developed tumors, while none of the three control animals developed tumors.

In another study, 15 mice were injected with mox1-transfected NIH 3T3 cells. Of the 15 mice injected, 14 showed large tumors within 17 days of injection, and tumors showed  
10 expression of mox1 mRNA. Histologically, the tumors resembled fibrosarcomas and were similar to ras-induced tumors. Thus, ras and mox1 were similarly potent in their ability to induce tumorigenicity of NIH 3T3 cells in athymic mice.

#### EXAMPLE 10

##### *Demonstration of the Role of Mox1 in Non-Cancerous Growth*

A role in normal growth was demonstrated in rat aortic vascular smooth-muscle cells by using antisense to rat  
20 mox1. Transfection with the antisense DNA resulted in a decrease in both superoxide generation and serum-dependent growth. Mox1 is therefore implicated in normal growth in this cell type.

#### EXAMPLE 11

##### *Expression of Human Mox1 Protein (SEQ ID NO:2) in a Baculovirus Expression System*

SEQ ID NO:2 was also expressed in insect cells using recombinant baculovirus. To establish the p65mox1  
30 expressing virus system, the mox1 gene (SEQ ID NO:1) was initially cloned into the pBacPAK8 vector (Clontech, Palo Alto, CA) and recombinant baculovirus was constructed using standard methods according to manufacturer's protocols. Briefly, PCR amplified mox1 DNA was cloned into the KpnI and EcoRI site of the vector. Primers used for PCR  
35 amplification were: 5'-CAA GGT ACC TCT TGA CCA TGG



GAA ACT-3' , SEQ ID NO:15, and 5'-ACG AAT TCA AGT  
AAA TTA CTG AAG ATA C-3' , SEQ ID NO:16. Sf9 insect  
cells ( $2 \times 10^6$  cells) were infected with 0.5 mg of linearized  
baculovirus DNA sold under the trademark BACULOGOLD®  
5 (PharMingen, San Diego, CA) and 5 mg pBacPAC8-p65mox1  
using Transfection Buffers A and B (PharMingen, San Diego,  
CA). After 5 days, the supernatants containing recombinant  
viruses were harvested and amplified by infecting fresh sf9 cells  
for 7 days. Amplification was carried out three times and the  
10 presence of the recombinant virus containing mox1 DNA was  
confirmed by PCR using the same primers. After three times  
amplification of viruses, plaque purification was carried out to  
obtain the high titer viruses. Approximately  $2 \times 10^8$  sf9 cells in  
15 agar plates were infected for 5 days with serial dilutions of  
virus and were dyed with neutral red for easy detection of virus  
plaques. Selected virus plaques were extracted and the presence  
of the human mox1 DNA was confirmed again by PCR.

### EXAMPLE 12

#### 20 *Cloning of a Rat Homolog of p65mox (SEQ ID NO:2)*

cDNA clones of p65mox from a rat aortic smooth  
muscle cell have been obtained. RT-PCR (reverse transcription  
polymerase chain reaction) was carried out as follows: first  
strand cDNA synthesis was performed using total RNA from rat  
25 aortic vascular smooth muscle cells, oligo dT primer and  
superscript II reverse transcriptase, and followed by incubation  
with RNase H. Degenerate PCR primers were designed to  
anneal to conserved areas in the coding regions of h-mox1 and  
gp91phox of human (X04011), mouse (U43384) and porcine  
30 (SSU02476) origin. Primers were: sense 5'-  
CCIGTITGTCGIAATCTGCTSTCCTT-3', SEQ ID NO:17 and  
antisense 5'-TCCCIGCAIAICCAAGTAGAARTAGATCTT-3',  
SEQ ID NO:18. A major PCR product of the expected 1.1 kb  
size was purified by agarose electrophoresis and used as  
35 template in a second PCR amplification reaction.

An aliquot of the RT-PCR product was blunt-



ended, ligated into a modified Litmus 29 vector and used to transform XL10 competent *E. coli*. Approximately 120 bacterial colonies were screened for the presence of a full-length insert by direct PCR using vector primers and Taq polymerase. Plasmids were purified from 25 positive colonies and mapped by digestion with *Bam* HI. Representative plasmids from each digestion pattern were partially sequenced. Five out of 25 clones contained non-specific amplification products and 20 contained identical inserts similar to human (h)-mox1. One of the latter clones was fully sequenced and found to be 83% identical to h-mox1 over 1060 nucleotides. A 1.1 kb probe was generated by PCR amplification of the insert of a rat mox1 clone with the degenerate primers described above and used to hybridize to a Northern blot of rat vascular smooth muscle cell RNA. A single band, migrating between 28S rRNA and 18S rRNA, indicated the presence of a message with a size compatible to that of human mox-1 (2.6 kb).

To obtain full-length rat mox1, 3' and 5' rapid amplification of cDNA ends (RACE) reactions were performed as describe above, using the gene-specific primers 5'-TTGGCACAGTCAGTGAGGATGTCTTC-3', SEQ ID NO:19 and 5'-CTGTTGGCTTCTACTGTAGCGTTCAAAGTT-3', SEQ ID NO:20 for 3' and 5' RACE, respectively. Single major 1.5 kb and 850 bp products were obtained for 3' and 5' RACE, respectively. These products were purified by agarose gel eletrophoresis and reamplified with Taq polymerase. Both products were cloned into the pCR 2.1 vector and used to transform electrocompetent XL1 blue *E. coli*. The RACE products were sequenced and new terminal primers were designed:

sense 5'-TTCTGAGTAGGTGTGCATTTGAGTGTCAATAAAGAC-3' (SEQ ID NO:43), and antisense 5'-TTTTCCGTCAAAATTATAACTTTTTATTTTCTTTTATAACACAT-3' (SEQ ID NO:44). PCR amplification of rat VSMC cDNA was performed using these primers.



A single 2.6 kb product was obtained, ligated into pCR 2.1 and used to transform electrocompetent XL1 blue *E. coli*. The insert was sequenced with 12 sense and 14 antisense primers. Its length is 2577 bp (including primer sequences), comprising a 1692 bp open reading frame, 127 bp 5' and 758 bp 3' untranslated regions. The presence of six in-frame stop codons in the 5' untranslated region suggests that the full length coding region has been obtained. Consensus polyadenylation sequences are present at nucleotides 2201 and 2550. Conceptual translation yields a 563 amino acid peptide, one residue shorter than the human deduced sequence. This new amino acid sequence is more similar to human mox1 SEQ ID NO:3 (82% identity) than to mouse gp91phox SEQ ID NO:38 (55% identity), suggesting that it is indeed rat mox1 (SEQ ID NO:21). This rat (r) homolog of p65mox protein is called r-p65mox or p65mox/rat.pep and is shown as SEQ ID NO:21. The nucleotide sequence encoding for r-p65mox is shown as SEQ ID NO:22.

20

### EXAMPLE 13

*Expression of rat (r)-p65mox mRNA in Vascular Smooth Muscle and Induction by Angiotensin II, Platelet-Derived Growth Factor (PDGF), and Phorbol Myristic Acid (PMA)*

25

30

35

Using the partial cDNA clone from rat, we have examined cultured rat aortic smooth muscle cells for expression of message for r-p65mox. We have observed the mRNA for r-p65mox in these cells. It has been previously reported (Griendling et al. (1994) *Circ. Res.* 74, 1141-1148; Fukui et al. (1997) *Circ. Res.* 80, 45-51; Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321) that *in vitro* or *in vivo* treatment with angiotensin II (AII) is a growth stimulus for vascular smooth muscle cells, and that AII induces increased superoxide generation in these cells. Platelet-derived growth factor (PDGF) and PMA are proliferative signals for vascular smooth muscle cells. We observed that the mRNA for r-p65mox was induced approximately 2-3 fold by angiotensin II (100 nM),



corresponding to the increased level of superoxide generation. Thus, the increased superoxide generation in these cells correlates with increased expression of the mRNA for this enzyme. The mRNA for r-p65mox also increased 2 or more fold in response to the growth stimulus PDGF (20 ng/ml), and 2-3 fold in response to PMA. Quantitation by densitometry revealed that rat mox1 message was induced nearly 4-fold at the 6 and 12 hour time points in response to PDGF, and about 2-fold at the 12 hour time point in response to AII. 28S RNA was used as a control for RNA recovery.

#### EXAMPLE 14

##### *Antibodies to Fragments of Human (h)-p65mox (SEQ ID NO:2)*

Polyclonal antibodies were raised in rabbits against the C-terminal half of h-p65mox (residues 233 through 564, SEQ ID NO:23) which is predicted to fold into a cytosolic domain containing FAD and the NADPH or NADH binding site. This domain was expressed in *E. coli* as an N-terminal GST-fusion protein and was purified on glutathione agarose by standard methods. Two anti-peptide antibodies were also made against h-p65mox (residues 243-256, referred to as Peptide A, SEQ ID NO:24) and h-65mox (residues 538-551, referred to as Peptide B, SEQ ID NO:25). Peptides were conjugated to keyhole limpet hemocyanin (KLH) using glutaraldehyde.

Antigens were injected into different rabbits initially in complete Freund's adjuvant, and were boosted 4 times with antigen in incomplete Freund's adjuvant at intervals of every three weeks. Approximately 0.5 mg to 1 mg of peptide was administered at each injection. Blood was drawn 1 week after each boost and a terminal bleed was carried out 2 weeks after the final boost. Antibodies to Peptide A and Peptide B were affinity purified by column chromatography through peptide A or peptide B conjugated to Affigel 15 (Bio-Rad, Richmond, CA). 10 mg of peptide was covalently crosslinked to 2 ml of Affigel 15 resin and the gel was washed with 20 ml of binding buffer (20 mM Hepes/NaOH, pH 7.0, 200 mM NaCl,



and 0.5 % Triton X-100). The remaining functional N-hydrosuccinimide was blocked with 100  $\mu$ l of 1 M ethanolamine. After washing with 20 ml of binding buffer, 5 ml of the antiserum was incubated with the pep A-conjugated Affigel 15 resin overnight at 4°C. Unbound protein was washed away with 20 ml of binding buffer. Elution of the antibodies from the gel was performed with 6 ml of elution buffer (100 mM glycine/HCl, pH 2.5, 200 mM NaCl, and 0.5% Triton X-100). The eluate was then neutralized by adding 0.9 ml of 1 M Tris/HCl, pH 8.0. The GST-fusion form of truncated p65mox1 protein (residues 233-566, SEQ ID NO:23) was expressed in *E. coli*. Samples (20  $\mu$ g each) were run on 12 % SDS-PAGE either before or 1 or 4 hours after induction with 100  $\mu$ M IPTG (isopropyl  $\beta$ -thiogalactoside).

The extracted proteins were subjected to immunoprobining with affinity purified antiserum to peptide A at a 1:1000 dilution. The detection of antigens was performed using an enhanced chemiluminescence kit (Amersham, Buckinghamshire, UK). The affinity purified antibody to mox1 (243-256, SEQ ID NO:24) was used at a dilution of 1:1000 in a Western blot in which a total of 10  $\mu$ g of protein was added to each lane. The major band observed at 4 hours after IPTG induction corresponded to the size of the GST-mox1 expressed in bacteria containing the pGEX-2T vector encoding the GST-mox1 fusion protein.

### Example 15

#### *Presence of an NAD(P)H Oxidase in Ras-Transformed Fibroblasts*

A superoxide-generating NADPH oxidase activity was detected in homogenates from NIH 3T3 cells, and this activity increased about 10-15 fold in Ras-transformed NIH 3T3 cells (Table 6). To establish the stable Ras-transformed cell lines, the DNA for human Ras encoding an activating mutation at amino acid number 12 (Valine, referred to as V12-Ras) was subcloned into BamH1 and EcoR1 sites of pCDNA3 vector



which has a neomycin resistance gene. V12-Ras in pCDNA3 and empty vector were transfected into NIH 3T3 cells using Lipofectamine Plus (Gibco).  $2 \times 10^6$  cells were maintained with DMEM containing 10% calf serum and transfected with 1 mg of DNA. After 2-days, cells were split and selected with the same medium but containing 1 mg/ml neomycin. Colonies surviving in selection media for 10 to 14 days were sub-cultured and characterized by immunoblot analysis using antibody against human-H-Ras.

The expression of Ras in cells transfected with pcDNA-3 vector alone or in three cell lines transfected with V12-Ras in the same vector was analyzed on a Western blot. The three cell lines were named V12-Ras-7, V12-Ras-4, and V12-Ras-8. The expression of V12-Ras varied widely among the three cell lines tested. The V12-Ras-4 cell line expressed the highest level of Ras followed by the V12-Ras-8 cell line. The V12-Ras-7 cell line expressed the lowest level of Ras.

Lysates from each of these lines were then prepared and tested for their ability to generate superoxide. For each cell line, cells were washed with cold HANKS' balanced salt solution (HBSS), collected by centrifugation, kept on dry-ice for more than 30 min, and disrupted by suspending in low salt buffer (LSB; 50 mM Tris/HCl, pH 7.5, 1 mM PMSF, and protease cocktail from Sigma) and passing through a syringe needle (18 gauge) ten times. Cell lysates were frozen in dry-ice immediately after determining the protein concentration.

Table 6 shows superoxide generation in the transfected cells measured using the lucigenin luminescence assay. For the assay, 5  $\mu$ g of cell lysates were incubated with the reaction mixture containing 10  $\mu$ M lucigenin (luminescent probe) and 100  $\mu$ M NADPH (substrate) in the presence or absence of 100  $\mu$ M arachidonate in the absence or presence of 100 U of superoxide dismutase (SOD) or 1  $\mu$ M diphenyleneiodonium (DPI). Luminescence of the reaction mixture was monitored for 0.5 second by LumiCounter (Packard) for four times at 3 second intervals. RLU in Table 5



refers to relative luminescence units.

As shown in Table 6, the luminescence was partially inhibited by superoxide dismutase indicating that the signal was due at least in part to the generation of superoxide. DPI, a known inhibitor of both neutrophil and non-neutrophil NADPH oxidase activities, completely inhibited activity. The generation of superoxide correlated with the expression of Ras in the three cell lines. Thus, oncogenic Ras appears to induce an NADPH-dependent-superoxide-generating activity that is similar to the activity catalyzed by p65mox1.

**Table 6**

RLU/5  $\mu$ g protein

		no additions	plus SOD	plus DPI
15	Vector Control (1)	465	154	48
	V12-Ras-7 (2)	1680	578	39
	V12-Ras-4 (3)	5975	2128	36
	V12-Ras-8 (4)	4883	2000	35

20

#### EXAMPLE 16

##### *Molecular Cloning of Another Rat mox1 cDNA Called Rat mox1B*

A rat cDNA library was screened in an effort to identify new rat mox sequences. The library was constructed in a ZAP express lambda phage vector (Stratagene, La Jolla, CA) using RNA isolated from rat vascular smooth muscle cells which had been exposed to 100 nM angiotensin II for 4 hours. The library was screened using standard blot hybridization techniques with the rat mox1 probe described previously. Fifteen individual clones were obtained that were characterized by PCR and restriction mapping. Two different types of clones were thus identified and representatives of each type were sequenced. A clone of the first type (representative of 13) was found to be similar to the previously identified rat mox1 and



was thus named rat *mox1B*. Clones of the second type (representative of 2) were incomplete rat *mox* sequences.

The length of the rat *mox1B* nucleotide sequence is 2619 bp and is listed as SEQ ID NO:41. The single longest  
5 1497 bp open reading frame encompasses nucleotides 362 to 1858. The presence of two in-frame stop codons in the 5' untranslated region at nucleotides 74 and 257 indicates that the full-length coding region has been isolated. Two putative polyadenylation sites are present at positions 2243 and 2592.  
10 Alignment of the rat *mox1* nucleotide sequence (SEQ ID NO:22) and the rat *mox1B* nucleotide sequence (SEQ ID NO:41) shows that the two nucleotides sequences are identical except at their 5' ends, suggesting that they may represent two alternatively spliced messages from the same gene. Sequence  
15 identity starts at nucleotides 269 and 311, for rat *mox1* and rat *mox1B*, respectively.

Conceptual translation of the rat *mox1B* nucleotide sequence (SEQ ID NO:41) yields a 499 amino acid sequence with a predicted molecular weight of 58 kDa. This amino acid  
20 sequence for rat *mox1B* protein is shown in SEQ ID NO:42. Alignment of the deduced amino acid sequences for rat *mox1* (SEQ ID NO:21) and rat *mox1B* (SEQ ID NO:42) indicates that rat *mox1B* is identical to rat *mox1A*, except for a missing stretch of 64 residues at the N-terminus. Therefore, rat *mox1B*  
25 appears to be a splicing variant derived from the same gene as rat *mox1*.

#### EXAMPLE 17

*Sequence Analysis and Cloning of the Human Mox2 cDNA (SEQ ID NO:3) Encoding for Production of the Human Mox2 Protein (SEQ ID NO:4)*  
30

Note that the *mox2* protein as described herein, was described in U.S. Provisional Application Serial No. 60/149,332 as *mox3*.

35 A blast search was carried out using the sequence of *mox1* as a query sequence. The sequence identified by this



search was a sequence present in the GenBank database that contains regions of homology with *mox1* and *gp91phox*. The GenBank sequence located in the search was a 90.6 kb sequenced region of human chromosome 6 (6q25.1-26) that was reported as a GenBank direct submission dated February 9, 1999 and given the Accession No. AL031773. Sequencing was carried out as part of the human genome sequencing project by S. Palmer, at Sanger Centre, in Hinxton, Cambridgeshire, UK. The GenBank sequence was reported as being similar to "Cytochrome B" and was not reported as having any homology or relation to a *mox* protein. The sequence contained a theoretical amino acid sequence that was derived by computer using an algorithm that predicted intron/exon boundaries and coding regions. This predicted region contained a 545 amino acid sequence that was 56% identical to *mox1* and 58% identical to *gp91phox*.

In the present invention, based on the GenBank genomic sequence and the homologies described above, several specific primers were designed and used to determine the tissue expression patterns of a novel *mox* protein, *mox2*, using Human Multiple Tissue PCR Panels (Clontech, Palo Alto, CA). The primers were as follows: Primer 1: 5'-CCTGACAGATGTATTTCACTACCCAG-3' (SEQ ID NO:49); Primer 2: 5'-GGATCGGAGTCACTCCCTTCGCTG-3' (SEQ ID NO:50); Primer 3: 5'-CTAGAAGCTCTCCTTGTGTAATAGA-3' (SEQ ID NO:51); Primer 4: 5'-ATGAACACCTCTGGGGTCAGCTGA-3' (SEQ ID NO:52). It was determined that *mox2* is expressed primarily in fetal tissues, with highest expression in fetal kidney, with expression also seen in fetal liver, fetal lung, fetal brain, fetal spleen and fetal thymus. Among 16 adult tissues tested, *mox2* expression was seen in brain, kidney, colon and lung, although levels of expression appeared to be very low.

Additionally, the 5' RACE (RACE = Rapid Amplification of cDNA Ends) and 3' RACE techniques were used to complete the sequence of the 5' and 3' regions of *mox2*.



(5' RACE kit and 3' RACE kit were from Clontech, Palo Alto, CA and are more fully described in Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85, 8998-9002. The 5' RACE and 3'-RACE techniques were carried out using a human fetal kidney library (Marathon-Ready cDNA library, Cat. #7423-1), using the following specific primers: 5'-RACE: Primer 4: 5'-ATGAACACCTCTGGGGTCAGCTGA-3' (SEQ ID NO:53); Primer 5: 5'-GTCCTCTGCAGCATTGTTCTCTTA-3' (SEQ ID NO:54); 3'-RACE: Primer 1: 5'-CCTGACAGATGTATTTCACTACCCAG-3' (SEQ ID NO:55); Primer 2: 5'-GGATCGGAGTCACTCCCTTCGCTG-3' (SEQ ID NO:56). The RACE procedures were successful in completing the 5' sequence and in confirming the 3' sequence. The complete coding sequence of *mox2* is shown in SEQ ID NO:2, while the predicted amino acid sequence of *mox2* is shown in SEQ ID NO:4.

In comparing the sequences of the present invention to the predicted coding regions of the GenBank sequence, the GenBank sequence did not contain a start codon, appeared to be missing approximately 45 base pairs at the N-terminus, and contained one other major difference in the predicted coding region which could have been due to inaccurate computer prediction of intron/exon boundaries.

25

#### EXAMPLE 18

*Sequence Analysis and Partial Cloning of the Human Duox2 cDNA (SEQ ID NO:47) Encoding for Production of the Human Duox2 Protein (SEQ ID NO:48)*

A partial cDNA clone of *duox2* was obtained as follows. A 535-base portion of an expressed sequence tag (EST zc92h03.rl; Genbank accession no. W52750) from human pancreatic islet was identified using the human gp91phox amino-acid sequence as a query in a Blast search. The bacterial strain #595758 containing the EST sequence zc92h03.rl in the pBluescript SK-vector was purchased from ATCC (Rockville, MD). The DNA inserted into the pBluescript SK-vector was



further sequenced using T7 and T3 vector promoters as well as sequence specific internal primers. The EST encoded 440 amino acids showing a 24.4% identity to gp91phox, including a stop codon corresponding to the C-terminus of gp91phox. 5'-RACE was carried out using mRNA obtained from human colon carcinoma cells (CaCo2) and the Marathon cDNA Amplification Kit (ClonTech, Palo Alto). The following gene-specific primers were used for this procedure: 5'-GAAGTGGTGGGAGGCGAAGACATA-3' (SEQ ID NO:26) and 5'-CCTGTCATACCTGGGACGGTCTGG-3' (SEQ ID NO:27).

The results of the 5'-RACE yielded an additional 2 kilobase of sequenced DNA but this region did not contain the start codon. To complete the sequence of the 5' and 3' regions of duox2, 5'-RACE and 3'-RACE were carried out using a human adult pancreas mRNA (Clontech, Palo Alto, CA) with the kit of 5' RACE System for Rapid Amplification of cDNA Ends version 2.0 (Gibco BRL, Gaithersburg, MD). PCR done using the following specific primers resulted in a total predicted amino acid sequence of about 1000 residues: 5'-RACE: Primer 3: 5'-GAGCACAGTGAGATGCCTGTTCAG-3' (SEQ ID NO:28); Primer 4: 5'-GGAAGGCAGCAGAGAGCAATGATG-3' (SEQ ID NO:29) (for nested PCR); 3'-RACE Primer 5: 5'-ACATCTGCGAGCGGCACTTCCAGA-3' (SEQ ID NO:30) Primer 6: 5'-AGCTCGTCAACAGGCAGGACCGAGC-3' (SEQ ID NO:31) (for nested PCR).

#### EXAMPLE 19

*Sequence Analysis and Cloning of the Human Duox1 cDNA (SEQ ID NO:45) Encoding for Production of the Human Duox1 Protein (SEQ ID NO:46)*

A cDNA clone of duox1 was obtained as follows. A homologous 357-base portion of an expressed sequence tag (EST nr80d12.s1; Genbank accession no. AA641653) from an invasive human prostate was identified by using the partial



duox2 predicted amino-acid sequence described above as a query in a Blast search. The bacterial strain #1441736 containing the EST sequence nr80d12.s1 in the pBluescript SK-vector was purchased from ATCC (Rockville, MD). The DNA inserted into the pBluescript SK-vector was further sequenced using T7 and T3 vector promoters as well as sequence specific internal primers. The EST insert encoded 673 amino acids with no start or stop codons present. Northern Blot analysis of duox1 indicated the gene was about 5.5 kilobase pairs. To complete the sequence of 5' and 3' regions of duox1, 5' RACE and 3'-RACE were carried out using a human adult lung mRNA (Clontech, Palo Alto, CA) with the kit of 5' RACE System for Rapid Amplification of cDNA Ends version 2.0 (Gibco BRL, Gaithersburg, MD). The RACE procedure was carried out using the following specific primers: 5'-RACE: Primer 5: 5'-GCAGTGCATCCACATCTTCAGCAC-3'(SEQ ID NO:32); Primer 6: 5'-GAGAGCTCTGGAGACACTTGAGTTC-3' (SEQ ID NO:33) (for nested PCR); 3'-RACE Primer 7: 5'-CATGTTCTCTCTGGCTGACAAG-3' (SEQ ID NO:34); Primer 8: 5'-CACAATAGCGAGCTCCGCTTCACGC-3' (SEQ ID NO:35) (for nested PCR). RACE procedures were successful in completing the 5' sequence and the 3' sequence of duox1. The open reading frame is approximately 4563 base pairs.

## EXAMPLE 20

### *Tissue Expression of Duox1 and Duox2*

Based on the duox1 sequence data, several specific primers were designed (Primer 1a: 5'-GCAGGACATCAACCCTGCACTCTC-3' (SEQ ID NO:36); Primer 2a: 5'-AATGACACTGTACTGGAGGCCACAG-3' (SEQ ID NO:57); Primer 3a: 5'-CTGCCATCTACCACACGGATCTGC-3' (SEQ ID NO:58); Primer 4a: 5'-CTTGCCATTCCAAAGCTTCCATGC-3' (SEQ ID NO:59) and used these to determine the tissue expression patterns of duox1 using Human Multiple Tissue PCR Panels



(Clontech, Palo Alto, CA). It was determined that duox1 is expressed primarily in lung, testis, placenta, prostate, pancreas, fetal heart, fetal kidney, fetal liver, fetal lung, fetal skeletal muscle and thymus, with highest expression in adult and fetal lung. Among 16 adult tissues and 8 fetal tissues tested, duox1 expression in brain, heart, kidney, colon, ovary, thymus, fetal brain and fetal spleen appeared to be low.

Two duox2 specific primers were also used to determine the tissue expression patterns of duox2 using Human Multiple Tissue PCR (polymerase chain reaction) Panels (Clontech, Palo Alto, CA). (Primer 1b: 5'-GTACAAGTCAGGACAGTGGGTGCG-3' (SEQ ID NO:60); Primer 2b: 5'-TGGATGATGTCAGCCAGCCACTCA-3' (SEQ ID NO:61)). Duox2 is expressed primarily in lung, pancreas, placenta, colon, prostate, testis and fetal tissues, with highest expression in adult lung and fetal tissues. Among 16 adult tissues and 8 fetal tissues tested, duox2 expression in brain, heart, kidney, liver, skeletal muscle, thymus and fetal brain appeared to be low.

## EXAMPLE 21

### *Role of Duox1 and Duox2 in Collagen Crosslinking*

To investigate a possible role for the human duox1 and duox2, the model organism *Caenorhabditis elegans* and a new reverse genetic tool, RNA interference (RNAi), were used to "knock out" the homologues of duox in this organism (Fire et al. (1998) *Nature* 391, 806-811). This technique involved injection of double stranded RNA encoding a segment of Ce-duox1 or Ce-duox2 into gonads of *C. elegans* N2 hermaphrodites. Injected worms were then allowed to lay eggs, and the harvested eggs were allowed to develop and the F1 progeny were scored for phenotypes. This procedure has been documented to "knock-out" the expression of the gene of interest (Fire et al. (1998) *Nature* 391, 806-811).

In the case of Ce-duox1 and Ce-duox2, the knockout animals resulted in a complex phenotype including



worms with large superficial blisters, short or "dumpy" worms, worms with locomotion disorders, and worms with retained eggs and/or larvae. Because of the high identity between Ce-duox1 and Ce-duox2, three different RNA constructs were predicted to knock out either both genes or Ce-duox2 alone. In all cases, essentially the same group of phenotypes was obtained. Most or all of these phenotypes had been described previously in *C. elegans* mutated in the collagen biosynthetic pathway. *C. elegans* has an extracellular structure known as the cuticle, a complex sheath composed largely of cross-linked collagen, which functions as the exoskeleton of the nematode. Cross-linking of collagen in nematodes occurs in part by cross-linking tyrosine residues, and peroxidases such as sea urchin ovoperoxidase and human myeloperoxidase have previously been shown to be capable of carrying out this reaction.

Based upon the similarities of the phenotypes obtained, the Ce-duox1/2 knockout worms were examined for the presence of dityrosine linkages, using an HPLC methodology (Andersen, S.O. (1966) *Acta Physiol. Scand.* 66, Suppl. 263-265; Abdelrahim et al. (1997) *J. Chromatogr. B Biomed. Sci. Appl.* 696, 175-182). It was determined that dityrosine linkages, while easily detected in the wild type worms, were almost completely lacking in the knockout worms. Thus, an inability to catalyze dityrosine cross-linking accounts for the phenotype of *C. elegans* failing to express Ce-duox1/2. These data support the concept that the duox enzymes in higher organisms can probably function in a similar manner to modulate the extracellular milieu, possibly the extracellular matrix and/or the basement membrane.

All patents, publications and abstracts cited above are incorporated herein by reference in their entirety. It should be understood that the foregoing relates only to preferred embodiments of the present invention and that numerous modifications or alterations may be made therein without



departing from the spirit and the scope of the present invention as defined in the following claims.



## CLAIMS

- 5 1. A protein capable of stimulating superoxide production, wherein the protein comprises mox or duox, a fragment thereof or a conservative substitution thereof.
- 10 2. The protein of Claim 1, wherein the protein, the fragment thereof, or the conservative substitution thereof comprises the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48, a fragment thereof, or a conservative substitution thereof.
- 15 3. A nucleotide sequence encoding for the protein, the fragment thereof or the conservative substitution thereof as recited in Claim 1.
- 20 4. The nucleotide sequence of Claim 3, wherein the nucleotide sequence comprises SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, or SEQ ID NO:47, a fragment thereof, or a conservative substitution thereof.
- 25 5. A vector, wherein the vector comprises a nucleotide sequence encoding for the protein, the fragment thereof or the conservative substitution thereof, as recited in Claim 1.
- 30 6. The vector of Claim 5 wherein the nucleotide sequence comprises SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, or SEQ ID NO:47, a fragment thereof, or a conservative substitution thereof.
- 35 7. A cell containing the vector of Claim 5.
8. A cell containing the vector of Claim 6.



9. An antibody, wherein the antibody is capable of binding to the protein, the fragment thereof, or the conservative substitution thereof, as recited in Claim 1.

5 10. The antibody of Claim 9, wherein the protein, the fragment thereof, or the conservative substitution thereof, has the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48, a fragment thereof, or a conservative substitution thereof.

10 11. A method of stimulating superoxide formation comprising administration, in vitro or in vivo, of a composition comprising the protein, the fragment thereof, or the conservative substitution thereof of Claim 1 in a  
15 pharmaceutically acceptable carrier.

20 12. The method of Claim 11, wherein the protein, the fragment thereof, or the conservative substitution thereof, comprises the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48, a fragment thereof, or a conservative substitution thereof.

25 13. A method of stimulating superoxide formation comprising administration, in vitro or in vivo, of a composition comprising the vector of Claim 5 in a pharmaceutically acceptable carrier.

30 14. A method of stimulating superoxide formation comprising administration, in vitro or in vivo, of a composition comprising the vector of Claim 6 in a pharmaceutically acceptable carrier.

35 15. A method for determining the activity of a drug comprising measuring the activity of the protein, the fragment thereof or the conservative substitution thereof, as recited in



Claim 1, to stimulate superoxide production following administration of the drug.

5

16. The method of Claim 15, wherein the protein, the fragment thereof or the conservative substitution thereof comprises the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48, a fragment thereof, or a conservative substitution thereof.

10



FIGURE 1(a)



154	F	A	R	K	R	I	K	N	P	E	G	G	L	Y	L	A	V	T	L	L	A	G	I	T	G	V	V	I	T	L	gp91phox.human		
155	F	V	R	Q	R	I	K	N	P	E	G	G	L	Y	V	A	V	T	R	L	A	G	I	T	G	V	V	I	T	L	gp91phox.bovine		
155	F	A	R	E	K	I	K	N	P	E	G	G	L	Y	V	A	V	T	R	L	A	G	I	T	G	I	V	I	T	L	gp91phox.mouse		
157	P	I	Q	S	R	-	-	-	N	T	T	V	E	Y	V	T	F	T	S	V	A	G	L	T	G	V	I	M	T	I	Mox1.human		
156	P	I	Q	S	P	-	-	-	N	V	T	V	M	Y	A	A	F	T	S	I	A	G	L	T	G	V	V	A	T	V	Mox1.rat		
156	P	V	R	T	F	P	T	N	T	T	T	E	L	R	T	I	-	-	A	G	V	T	G	L	V	I	S	L	mox2.human				
169	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	M	K	G	T	D	G	W	T	G	V	T	M	V	cytb558/arabidopsis.pep		
162	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	V	K	G	V	E	G	I	T	G	V	I	M	V	cytb558.rice.pep		
184	C	L	I	L	I	I	T	S	S	T	K	T	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	F	gp91phox.human	
185	C	L	I	L	I	I	T	S	S	T	K	T	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	gp91phox.bovine
185	C	L	I	L	I	I	T	S	S	T	K	T	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	gp91phox.mouse
184	A	L	I	L	M	V	T	S	A	T	E	F	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	Mox1.human
183	A	L	V	L	M	V	T	S	A	M	E	F	I	R	R	N	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	Mox1.rat
183	A	L	V	L	I	M	T	S	S	T	E	F	I	R	Q	A	S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Y	mox2.human
183	L	M	L	V	A	Y	V	L	A	Q	S	W	F	R	R	N	R	A	N	L	P	K	S	L	K	R	L	T	G	F	cytb558/arabidopsis.pep		
176	C	M	I	I	A	F	T	L	A	T	R	W	F	R	R	S	L	V	K	L	P	R	P	F	D	K	L	T	G	F	cytb558.rice.pep		
202	E	V	F	W	Y	T	H	H	L	F	V	I	F	F	I	G	L	A	I	H	G	A	E	R	I	V	R	G	Q	T	gp91phox.human		
203	E	V	F	W	Y	T	H	H	L	F	V	I	F	F	I	G	L	A	I	H	G	A	Q	R	I	V	R	G	Q	T	gp91phox.bovine		
203	E	V	F	W	Y	T	H	H	L	F	V	I	F	F	I	G	L	A	I	H	G	A	E	R	I	V	R	G	Q	T	gp91phox.mouse		
202	E	V	F	W	Y	T	H	H	L	F	I	F	Y	I	L	G	L	G	I	H	G	I	G	G	I	V	R	G	Q	T	Mox1.human		
201	E	L	F	W	Y	T	H	H	L	F	I	I	Y	I	I	C	L	G	I	H	G	L	G	G	I	V	R	G	Q	T	Mox1.rat		
201	E	L	F	W	Y	T	H	H	V	F	I	V	F	E	L	S	L	A	I	H	G	T	G	R	I	V	R	G	Q	T	mox2.human		
213	N	A	F	W	Y	S	H	H	L	F	V	I	V	Y	V	L	L	I	V	H	G	-	-	-	-	-	-	-	-	-	-	cytb558/arabidopsis.pep	
206	N	A	F	W	Y	S	H	H	L	F	I	I	V	Y	I	A	L	I	V	H	G	-	-	-	-	-	-	-	-	-	-	cytb558.rice.pep	
232	A	E	S	L	A	V	H	N	I	T	V	C	E	Q	K	I	S	E	W	G	K	I	K	-	E	C	P	I	P	O	gp91phox.human		
233	A	E	S	L	L	K	H	Q	P	R	N	C	Y	Q	N	I	S	Q	W	G	K	I	E	-	N	C	P	I	P	E	gp91phox.bovine		
233	A	E	S	L	E	H	N	L	D	I	C	A	D	K	I	E	E	W	G	K	I	K	-	E	C	P	V	P	K	gp91phox.mouse			
232	E	E	S	M	N	E	S	H	P	R	K	C	A	E	S	F	E	M	W	D	D	R	D	S	H	C	R	R	P	K	Mox1.human		
231	E	E	S	M	S	E	S	H	P	R	N	C	S	Y	S	F	H	E	W	D	K	Y	E	R	S	C	R	S	P	H	Mox1.rat		
231	Q	D	S	L	S	L	H	N	I	T	F	C	R	D	R	Y	A	E	W	Q	T	V	A	-	Q	C	P	V	P	O	mox2.human		
234	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	cytb558/arabidopsis.pep	
227	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	cytb558.rice.pep	
261	F	A	G	N	P	P	M	T	W	K	W	I	V	G	P	M	F	L	Y	L	C	E	R	L	V	R	F	W	R	S	gp91phox.human		
262	F	S	G	N	P	P	M	T	W	K	W	I	V	G	P	M	F	L	Y	L	C	E	R	L	V	R	F	W	R	S	gp91phox.bovine		
262	F	A	G	N	P	P	M	T	W	K	W	I	V	G	P	M	F	L	Y	L	C	E	R	L	V	R	F	W	R	S	gp91phox.mouse		
262	F	E	G	H	P	P	E	S	W	K	W	I	L	A	P	V	I	L	Y	I	C	E	R	I	L	R	F	Y	R	S	Mox1.human		
261	F	V	G	Q	P	P	E	S	W	K	W	I	L	A	P	I	A	F	Y	I	F	E	R	I	L	R	F	Y	R	S	Mox1.rat		
260	F	S	G	K	E	P	S	A	W	K	W	I	L	G	P	V	V	L	Y	A	C	E	R	I	I	R	F	W	R	F	mox2.human		
243	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	cytb558/arabidopsis.pep	
236	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	cytb558.rice.pep	
291	Q	Q	K	V	V	-	-	-	I	T	K	V	V	T	H	P	F	K	T	I	E	L	Q	M	K	K	-	K	G	F	K	M	gp91phox.human
292	Q	Q	K	V	V	-	-	-	I	T	K	V	V	T	H	P	F	K	T	I	E	L	Q	M	K	K	-	K	G	F	K	M	gp91phox.bovine
292	Q	Q	K	V	V	-	-	-	I	T	K	V	V	T	H	P	F	K	T	I	E	L	Q	M	K	K	-	K	G	F	K	M	gp91phox.mouse
292	Q	Q	K	V	V	-	-	-	I	T	K	V	V	M	H	P	S	K	V	L	E	L	Q	M	N	K	-	R	G	F	S	M	Mox1.human
291	R	Q	K	V	V	-	-	-	I	T	K	V	V	M	H	P	C	K	V	L	E	L	Q	M	R	K	-	R	G	F	T	M	Mox1.rat
290	Q	Q	E	V	V	-	-	-	I	T	K	V	V	S	H	P	S	G	V	L	E	L	H	M	K	K	-	R	G	F	K	M	mox2.human
270	G	A	K	A	V	K	V	L	K	V	A	V	Y	P	G	N	V	L	S	L	Y	M	S	K	P	K	G	F	K	Y	cytb558/arabidopsis.pep		
263	G	S	Y	S	V	R	L	L	K	V	A	I	Y	P	G	N	V	L	T	L	O	M	S	K	P	P	T	E	R	Y	cytb558.rice.pep		

FIGURE 1(b)



3/8

319	E V G Q Y I F V K C P K V S K L E W H P F T L T S A P E E D	gp91phox.human
320	E V G Q Y I F V K C P V V S K L E W H P F T L T S A P E E D	gp91phox.bovine
320	E V G Q Y I F V K C P K V S K L E W H P F T L T S A P E E D	gp91phox.mouse
320	E V G Q Y I F V N C P S I S L L E W H P F T L T S A P E E D	Mox1.human
319	G I G Q Y I F V N C P S I S F L E W H P F T L T S A P E E E	Mox1.rat
318	A P G Q Y I L V Q C P A I S S L E W H P F T L T S A P Q E D	mox2.human
300	T S G Q Y I Y I N C S D V S P L Q W H P F S I T S A S G D D	cytb558/arabidopsis.pep
293	K S G Q Y M F V Q C P A V S P F E W H P F S I T S A P G D D	cytb558.rice.pep
349	F F S I H I R I V G D W T E G L F N A C G - - C - - - - -	gp91phox.human
350	F F S I H I R I V G D W T E G L F K A C G - - C - - - - -	gp91phox.bovine
350	F F S I H I R I V G D W T E G L F N A C G - - C - - - - -	gp91phox.mouse
350	F F S I H I R A A G D W T E N L I R A F E - - - - -	Mox1.human
349	F F S I H I R A A G D W T E N L I R T F E - - - - -	Mox1.rat
348	F F S V H I R A A G D W T A A L L E A F G - - A - - - - -	mox2.human
330	Y L S V H I R T L G D W T S Q L K S L Y S K V C Q L P S T S	cytb558/arabidopsis.pep
323	Y L S I H V R Q L G D W T R E L K R V F A A A C E P P A G G	cytb558.rice.pep
371	- - - - - D K Q E F Q D A W K L P K I A V D G P F G T A	gp91phox.human
372	- - - - - D K Q E F Q D A W K L P K I A V D G P F G T A	gp91phox.bovine
372	- - - - - D K Q E F Q D A W K L P K I A V D G P F G T A	gp91phox.mouse
371	- - - - - Q Q Y S P I P R I E V D G P F G T A	Mox1.human
370	- - - - - Q Q H S P M P R I E V D G P F G T V	Mox1.rat
370	- - - - - E G Q A L Q E P W S L P R L A V D G P F G T A	mox2.human
360	Q S G L F I A D I G Q A N N I T R F P R L L I D G P Y G A P	cytb558/arabidopsis.pep
353	K S G L L R A D - E T T K K I - - L P K L L I D G P Y G S P	cytb558.rice.pep
394	S E D V F S Y E V V M L V G A G I G V T P F A S I L K S V W	gp91phox.human
395	S E D V F S Y E V V M L V G A G I G V T P F A S I L K S V W	gp91phox.bovine
395	S E D V F S Y E V V M L V G A G I G V T P F A S I L K S V W	gp91phox.mouse
389	S E D V F Q Y E V A V L V G A G I G V T P F A S I L K S I W	Mox1.human
388	S E D V F Q Y E V A V L V G A G I G V T P F A S F L K S I W	Mox1.rat
393	L T D V F H Y P V C V C V A A G I G V T P F A A L L K S I W	mox2.human
390	A Q D Y R N Y D V L L L V G L G I G A T P L I S I I R D V L	cytb558/arabidopsis.pep
380	A Q D Y S K Y D V L L L V G L G I G A T P F I S I L K D L L	cytb558.rice.pep
424	Y K Y C N N A T N - - - - -	gp91phox.human
425	Y K Y C N K A P N - - - - -	gp91phox.bovine
425	Y K Y C D N A T S - - - - -	gp91phox.mouse
419	Y K F Q C A D H N - - - - -	Mox1.human
418	Y K F Q R A H N K - - - - -	Mox1.rat
423	Y K C S E A Q T P - - - - -	mox2.human
420	N N I - - - - - K N Q N S I E R G	cytb558/arabidopsis.pep
410	N N I I K M E E E E D A S T D L Y P P M G R N N P H V D L G	cytb558.rice.pep
433	- - - - - L K L K K I Y F Y W L C R D T H A F	gp91phox.human
434	- - - - - L R L K K I Y F Y W L C R D T H A F	gp91phox.bovine
434	- - - - - L K L K K I Y F Y W L C R D T H A F	gp91phox.mouse
428	- - - - - L K T K K I Y F Y W I C R E T G A F	Mox1.human
427	- - - - - L K T Q K I Y F Y W I C R E T G A F	Mox1.rat
432	- - - - - L K L S K V Y F Y W I C R D A R A F	mox2.human
432	T N Q H I - - - K N Y V A T K R A Y F Y W V T R E Q G S L	cytb558/arabidopsis.pep
440	T L M T I T S R P K K I L K T T N A Y F Y W V T R E Q G S F	cytb558.rice.pep

FIGURE 1(c)



4/8

451	EWFADLLQLLESQMQRNNAGFLSY	--NIY	gp91phox.human
452	EWFADLLQLLETQMQRNNAGFLSY	--NIC	gp91phox.bovine
452	EWFADLLQLLETQMQRNNAGFLSY	--NIY	gp91phox.mouse
446	SWFNLLTSLLEQEMELGKVGFLNY	--RLF	Mox1.human
445	AWFNLLTSLLEQEMDELGKPDFLNY	--RLF	Mox1.rat
450	EWFADLLTSLLETRMSQGGKTHFLSY	--HIF	mox2.human
458	EWFSEVMNEV-AEYDSEGMIELHNY	CTSVY	cytb558/arabidopsis.pep
470	DWFKGVMMNEI-ADLDQRNIIEMHNY	LTSVY	cytb558.rice.pep
479	LTG--WDESQANHFHAVHHDEEK	-DVITGLK	gp91phox.human
480	LTG--WDESQASHFAMHHDEEK	-DVITGLK	gp91phox.bovine
480	LTG--WDESQANHFHAVHHDEEK	-DVITGLK	gp91phox.mouse
474	LTG--WDSNIIVGHAAALNFDKAT	-DIVTGLK	Mox1.human
473	LTG--WDSNIAGHAAALNFDKAT	-DIVTGLK	Mox1.rat
478	LTG--WDENOALHIALHWDENT	-DVITGLK	mox2.human
487	EEDARSAALITMLQSLHHAHAKSGID	IVSGTR	cytb558/arabidopsis.pep
499	EEDARSAALITMLQALNHAKNGVD	IVSGTK	cytb558.rice.pep
506	QKTLYGRPNWDNEFKTIASQHPNT	RIGVFL	gp91phox.human
507	QKTLYGRPNWDNEFKTIASQHPNT	RIGVFL	gp91phox.bovine
507	QKTLYGRPNWDNEFKTIASQHPNT	RIGVFL	gp91phox.mouse
501	QKTSFGRPMWDNEFSSTIATSHPK	SVVGFL	Mox1.human
500	QKTSFGRPMWDNEFSSTIATSHPK	SVVGFL	Mox1.rat
505	QKTFYGRPNWNNEFKQIAYNHPS	SSIGVFF	mox2.human
517	VRTHFARPNWRSVFKHVAVNHVNQ	RVGVFY	cytb558/arabidopsis.pep
529	VRTHFARPNWRKVLISKISSKHPYA	KIGVFY	cytb558.rice.pep
536	CGPEALAEATLSKQSSISNS	ESGPRGVHFIFN	gp91phox.human
537	CGPEALAEATLSKQSSISNS	DSGPRGVHFIFN	gp91phox.bovine
537	CGPEALAEATLSKQSSISNS	ESGPRGVHFIFN	gp91phox.mouse
531	CGPRTLAKSLRKCCCHRYSSLDPRK	VQFYFN	Mox1.human
530	CGPRTLAKSLRKCCCHRYSSLDPRK	VQFYFN	Mox1.rat
535	CGPKALSLRTLQKMCCHLYSSADPR	GVHFYFN	mox2.human
547	CGNTCIIIGELKRLAQDFSRKT	--TTKFEFH	cytb558/arabidopsis.pep
559	CGAPVLAQELSKLCHFEFNGKC	--TTKFEFH	cytb558.rice.pep
566	KENF		gp91phox.human
567	KENF		gp91phox.bovine
567	KENF		gp91phox.mouse
561	KEN-F		Mox1.human
560	KETF		Mox1.rat
565	KESF		mox2.human
575	KENF		cytb558/arabidopsis.pep
587	KEHF		cytb558.rice.pep

Decoration 'Decoration #1': Box residues that match the Consensus exactly.

FIGURE 1(d)



5/8




FIGURE 2



6/8




FIGURE 3



7/8




FIGURE 4



8/8




FIGURE 5



## SEQUENCE LISTING

&lt;110&gt; Emory University

&lt;120&gt; Novel Mitogenic Regulators

&lt;130&gt; 05501-0103wp

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 61

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 2609

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (207)..(1901)

&lt;400&gt; 1

```

gctgatagca cagttctgtc cagagaagga aggcggaata aacttattca ttcccaggaa 60
ctcttggggg aggtgtgtgt ttttcacatc ttaaaggctc acagaccctg cgctggacaa 120
atgttccatt cctgaaggac ctctccagaa tccggattgc tgaatcttcc ctgttgcccta 180
gaagggtccc aaaccacctc ttgaca atg gga aac tgg gtg gtt aac cac tgg 233
                               Met Gly Asn Trp Val Val Asn His Trp
                               1             5

ttt tca gtt ttg ttt ctg gtt gtt tgg tta ggg ctg aat gtt ttc ctg 281
Phe Ser Val Leu Phe Leu Val Val Trp Leu Gly Leu Asn Val Phe Leu
 10             15             20             25

ttt gtg gat gcc ttc ctg aaa tat gag aag gcc gac aaa tac tac tac 329
Phe Val Asp Ala Phe Leu Lys Tyr Glu Lys Ala Asp Lys Tyr Tyr Tyr
      30             35             40

aca aga aaa atc ctt ggg tca aca ttg gcc tgt gcc cga gcg tct gct 377
Thr Arg Lys Ile Leu Gly Ser Thr Leu Ala Cys Ala Arg Ala Ser Ala
      45             50             55

ctc tgc ttg aat ttt aac agc acg ctg atc ctg ctt cct gtg tgt cgc 425

```



Leu Cys Leu Asn Phe Asn Ser Thr Leu Ile Leu Leu Pro Val Cys Arg	
60 65 70	
aat ctg ctg tcc ttc ctg agg ggc acc tgc tca ttt tgc agc cgc aca	473
Asn Leu Leu Ser Phe Leu Arg Gly Thr Cys Ser Phe Cys Ser Arg Thr	
75 80 85	
ctg aga aag caa ttg gat cac aac ctc acc ttc cac aag ctg gtg gcc	521
Leu Arg Lys Gln Leu Asp His Asn Leu Thr Phe His Lys Leu Val Ala	
90 95 100 105	
tat atg atc tgc cta cat aca gct att cac atc att gca cac ctg ttt	569
Tyr Met Ile Cys Leu His Thr Ala Ile His Ile Ile Ala His Leu Phe	
110 115 120	
aac ttt gac tgc tat agc aga agc cga cag gcc aca gat ggc tcc ctt	617
Asn Phe Asp Cys Tyr Ser Arg Ser Arg Gln Ala Thr Asp Gly Ser Leu	
125 130 135	
gcc tcc att ctc tcc agc cta tct cat gat gag aaa aag ggg ggt tct	665
Ala Ser Ile Leu Ser Ser Leu Ser His Asp Glu Lys Lys Gly Gly Ser	
140 145 150	
tgg cta aat ccc atc cag tcc cga aac acg aca gtg gag tat gtg aca	713
Trp Leu Asn Pro Ile Gln Ser Arg Asn Thr Thr Val Glu Tyr Val Thr	
155 160 165	
ttc acc agc gtt gct ggt ctc act gga gtg atc atg aca ata gcc ttg	761
Phe Thr Ser Val Ala Gly Leu Thr Gly Val Ile Met Thr Ile Ala Leu	
170 175 180 185	
att ctc atg gta act tca gct act gag ttc atc cgg agg agt tat ttt	809
Ile Leu Met Val Thr Ser Ala Thr Glu Phe Ile Arg Arg Ser Tyr Phe	
190 195 200	
gaa gtc ttc tgg tat act cac cac ctt ttt atc ttc tat atc ctt ggc	857
Glu Val Phe Trp Tyr Thr His His Leu Phe Ile Phe Tyr Ile Leu Gly	
205 210 215	
tta ggg att cac ggc att ggt gga att gtc cgg ggt caa aca gag gag	905
Leu Gly Ile His Gly Ile Gly Gly Ile Val Arg Gly Gln Thr Glu Glu	
220 225 230	
agc atg aat gag agt cat cct cgc aag tgt gca gag tct ttt gag atg	953
Ser Met Asn Glu Ser His Pro Arg Lys Cys Ala Glu Ser Phe Glu Met	
235 240 245	
tgg gat gat cgt gac tcc cac tgt agg cgc cct aag ttt gaa ggg cat	1001



3



Thr Gly Ala Phe Ser Trp Phe Asn Asn Leu Leu Thr Ser Leu Glu Gln  
 445 450 455  
 gag atg gag gaa tta ggc aaa gtg ggt ttt cta aac tac cgt ctc ttc 1625  
 Glu Met Glu Glu Leu Gly Lys Val Gly Phe Leu Asn Tyr Arg Leu Phe  
 460 465 470  
 ctc acc gga tgg gac agc aat att gtt ggt cat gca gca tta aac ttt 1673  
 Leu Thr Gly Trp Asp Ser Asn Ile Val Gly His Ala Ala Leu Asn Phe  
 475 480 485  
 gac aag gcc act gac atc gtg aca ggt ctg aaa cag aaa acc tcc ttt 1721  
 Asp Lys Ala Thr Asp Ile Val Thr Gly Leu Lys Gln Lys Thr Ser Phe  
 490 495 500 505  
 ggg aga cca atg tgg gac aat gag ttt tct aca ata gct acc tcc cac 1769  
 Gly Arg Pro Met Trp Asp Asn Glu Phe Ser Thr Ile Ala Thr Ser His  
 510 515 520  
 ccc aag tct gta gtg gga gtt ttc tta tgt ggc cct cgg act ttg gca 1817  
 Pro Lys Ser Val Val Gly Val Phe Leu Cys Gly Pro Arg Thr Leu Ala  
 525 530 535  
 aag agc ctg cgc aaa tgc tgt cac cga tat tcc agt ctg gat cct aga 1865  
 Lys Ser Leu Arg Lys Cys Cys His Arg Tyr Ser Ser Leu Asp Pro Arg  
 540 545 550  
 aag gtt caa ttc tac ttc aac aaa gaa aat ttt tga gttataggaa 1911  
 Lys Val Gln Phe Tyr Phe Asn Lys Glu Asn Phe  
 555 560 565  
 taaggacggt aatctgcatt ttgtctcttt gtatcttcag taattgagtt ataggaataa 1971  
 ggacggtaat ctgcattttg tctctttgta tcttcagtaa tttacttggt ctcttcaggt 2031  
 ttgancagtc actttaggat aagaatgtgc ctctcaagcc ttgactccct ggtattcttt 2091  
 ttttgattgc attcaacttc gttacttgag cttcagcaac ttaagaactt ctgaagttct 2151  
 taaagttctg aanttcttaa agcccatgga tcctttctca gaaaaataac tgtaaattctt 2211  
 tctggacagc catgactgta gcaaggcttg atagcagaag tttgggtggt canaattata 2271  
 caactaatcc caggtgattt tatcaattcc agtgttacca tctcctgagt tttgggtttgt 2331  
 aatcttttgt cctccacc cccacagaag attttaagta gggtgacttt ttaaataaaa 2391  
 atttattgaa taattaatga taaaacataa taataaacat aaataataaa caaaattacc 2451



gagaacccca tcccatata acaccaacag tgtacatgtt tactgtcact tttgatatgg 2511

tttatccagt gtgaacagca atttattatt tttgctcatc aaaaaataaa ggattttttt 2571

tcacttgaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2609

<210> 2

<211> 564

<212> PRT

<213> Homo sapiens

<400> 2

Met Gly Asn Trp Val Val Asn His Trp Phe Ser Val Leu Phe Leu Val  
1 5 10 15

Val Trp Leu Gly Leu Asn Val Phe Leu Phe Val Asp Ala Phe Leu Lys  
20 25 30

Tyr Glu Lys Ala Asp Lys Tyr Tyr Tyr Thr Arg Lys Ile Leu Gly Ser  
35 40 45

Thr Leu Ala Cys Ala Arg Ala Ser Ala Leu Cys Leu Asn Phe Asn Ser  
50 55 60

Thr Leu Ile Leu Leu Pro Val Cys Arg Asn Leu Leu Ser Phe Leu Arg  
65 70 75 80

Gly Thr Cys Ser Phe Cys Ser Arg Thr Leu Arg Lys Gln Leu Asp His  
85 90 95

Asn Leu Thr Phe His Lys Leu Val Ala Tyr Met Ile Cys Leu His Thr  
100 105 110

Ala Ile His Ile Ile Ala His Leu Phe Asn Phe Asp Cys Tyr Ser Arg  
115 120 125

Ser Arg Gln Ala Thr Asp Gly Ser Leu Ala Ser Ile Leu Ser Ser Leu  
130 135 140

Ser His Asp Glu Lys Lys Gly Gly Ser Trp Leu Asn Pro Ile Gln Ser  
145 150 155 160

Arg Asn Thr Thr Val Glu Tyr Val Thr Phe Thr Ser Val Ala Gly Leu  
165 170 175

Thr Gly Val Ile Met Thr Ile Ala Leu Ile Leu Met Val Thr Ser Ala



180	185	190
Thr Glu Phe Ile Arg Arg Ser Tyr Phe Glu Val Phe Trp Tyr Thr His 195	200	205
His Leu Phe Ile Phe Tyr Ile Leu Gly Leu Gly Ile His Gly Ile Gly 210	215	220
Gly Ile Val Arg Gly Gln Thr Glu Glu Ser Met Asn Glu Ser His Pro 225	230	235 240
Arg Lys Cys Ala Glu Ser Phe Glu Met Trp Asp Asp Arg Asp Ser His 245	250	255
Cys Arg Arg Pro Lys Phe Glu Gly His Pro Pro Glu Ser Trp Lys Trp 260	265	270
Ile Leu Ala Pro Val Ile Leu Tyr Ile Cys Glu Arg Ile Leu Arg Phe 275	280	285
Tyr Arg Ser Gln Gln Lys Val Val Ile Thr Lys Val Val Met His Pro 290	295	300
Ser Lys Val Leu Glu Leu Gln Met Asn Lys Arg Gly Phe Ser Met Glu 305	310	315 320
Val Gly Gln Tyr Ile Phe Val Asn Cys Pro Ser Ile Ser Leu Leu Glu 325	330	335
Trp His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Asp Phe Phe Ser 340	345	350
Ile His Ile Arg Ala Ala Gly Asp Trp Thr Glu Asn Leu Ile Arg Ala 355	360	365
Phe Glu Gln Gln Tyr Ser Pro Ile Pro Arg Ile Glu Val Asp Gly Pro 370	375	380
Phe Gly Thr Ala Ser Glu Asp Val Phe Gln Tyr Glu Val Ala Val Leu 385	390	395 400
Val Gly Ala Gly Ile Gly Val Thr Pro Phe Ala Ser Ile Leu Lys Ser 405	410	415
Ile Trp Tyr Lys Phe Gln Cys Ala Asp His Asn Leu Lys Thr Lys Lys 420	425	430
Ile Tyr Phe Tyr Trp Ile Cys Arg Glu Thr Gly Ala Phe Ser Trp Phe		



```
<210> 3
<211> 2044
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> CDS
<222> (104)..(1810)
```

```

<400> 3
caaagacaaa ataatttact agggaagccc ttactaacga cccaacatcc agacacaggt 60

gaggggagaag aaatttcctg acagccgaag agcaacaagt atc atg atg ggg tgc 115
                                         Met Met Gly Cys
                                         1

tgg att ttg aat gag ggt ctc tcc acc ata tta gta ctc tca tgg ctg 163
Trp Ile Leu Asn Glu Gly Leu Ser Thr Ile Leu Val Leu Ser Trp Leu
  5              10              15              20

gga ata aat ttt tat ctg ttt att gac acg ttc tac tgg tat gaa gag 211
Gly Ile Asn Phe Tyr Leu Phe Ile Asp Thr Phe Tyr Trp Tyr Glu Glu

```



	25	30	35	
gag gag tct ttc cat tac aca cga gtt att ttg ggt tca aca ctg gct				259
Glu Glu Ser Phe His Tyr Thr Arg Val Ile Leu Gly Ser Thr Leu Ala				
	40	45	50	
tgg gca cga gca tcc gca ctg tgc ctg aat ttt aac tgc atg cta att				307
Trp Ala Arg Ala Ser Ala Leu Cys Leu Asn Phe Asn Cys Met Leu Ile				
	55	60	65	
cta ata cct gtc agt cga aac ctt att tca ttc ata aga gga aca agt				355
Leu Ile Pro Val Ser Arg Asn Leu Ile Ser Phe Ile Arg Gly Thr Ser				
	70	75	80	
att tgc tgc aga gga ccg tgg agg agg caa tta gac aaa aac ctc aga				403
Ile Cys Cys Arg Gly Pro Trp Arg Arg Gln Leu Asp Lys Asn Leu Arg				
	85	90	95	100
ttt cac aaa ctg gtc gcc tat ggg ata gct gtt aat gca acc atc cac				451
Phe His Lys Leu Val Ala Tyr Gly Ile Ala Val Asn Ala Thr Ile His				
	105	110	115	
atc gtg gcg cat ttc ttc aac ctg gaa cgc tac cac tgg agc cag tcc				499
Ile Val Ala His Phe Phe Asn Leu Glu Arg Tyr His Trp Ser Gln Ser				
	120	125	130	
gag gag gcc cag gga ctt ctg gcc gca ctt tcc aag ctg ggc aac acc				547
Glu Glu Ala Gln Gly Leu Leu Ala Ala Leu Ser Lys Leu Gly Asn Thr				
	135	140	145	
cct aac gag agc tac ctc aac cct gtc cgg acc ttc ccc aca aac aca				595
Pro Asn Glu Ser Tyr Leu Asn Pro Val Arg Thr Phe Pro Thr Asn Thr				
	150	155	160	
acc act gaa ttg cta agg aca ata gca ggc gtc acc ggt ctg gtg atc				643
Thr Thr Glu Leu Leu Arg Thr Ile Ala Gly Val Thr Gly Leu Val Ile				
	165	170	175	180
tct ctg gct tta gtc ttg atc atg acc tcg tca act gag ttc atc aga				691
Ser Leu Ala Leu Val Leu Ile Met Thr Ser Ser Thr Glu Phe Ile Arg				
	185	190	195	
cag gcc tcc tat gag ttg ttc tgg tac aca cac cat gtt ttc atc gtc				739
Gln Ala Ser Tyr Glu Leu Phe Trp Tyr Thr His His Val Phe Ile Val				
	200	205	210	
ttc ttt ctc agc ctg gcc atc cat ggg acg ggt cgg att gtt cga ggc				787
Phe Phe Leu Ser Leu Ala Ile His Gly Thr Gly Arg Ile Val Arg Gly				



215	220	225	
caa acc caa gac agt ctc tct ctg cac aac atc acc ttc tgt aga gac			835
Gln Thr Gln Asp Ser Leu Ser Leu His Asn Ile Thr Phe Cys Arg Asp			
230	235	240	
cgc tat gca gaa tgg cag aca gtg gcc caa tgc ccc gtg cct caa ttt			883
Arg Tyr Ala Glu Trp Gln Thr Val Ala Gln Cys Pro Val Pro Gln Phe			
245	250	255	260
tct ggc aag gaa ccc tcg gct tgg aaa tgg att tta ggc cct gtg gtc			931
Ser Gly Lys Glu Pro Ser Ala Trp Lys Trp Ile Leu Gly Pro Val Val			
265	270	275	
ttg tat gca tgt gaa aga ata att agg ttc tgg cga ttt caa caa gaa			979
Leu Tyr Ala Cys Glu Arg Ile Ile Arg Phe Trp Arg Phe Gln Gln Glu			
280	285	290	
gtt gtc att acc aag gtg gta agc cac ccc tct gga gtc ctg gaa ctt			1027
Val Val Ile Thr Lys Val Val Ser His Pro Ser Gly Val Leu Glu Leu			
295	300	305	
cac atg aaa aag cgt ggc ttt aaa atg gcg cca ggg cag tac atc ttg			1075
His Met Lys Lys Arg Gly Phe Lys Met Ala Pro Gly Gln Tyr Ile Leu			
310	315	320	
gtg cag tgc cca gcc ata tct tcg ctg gag tgg cac ccc ttc acc ctt			1123
Val Gln Cys Pro Ala Ile Ser Ser Leu Glu Trp His Pro Phe Thr Leu			
325	330	335	340
acc tct gcc ccc cag gaa gac ttt ttc agc gtg cac atc cgg gca gca			1171
Thr Ser Ala Pro Gln Glu Asp Phe Phe Ser Val His Ile Arg Ala Ala			
345	350	355	
gga gac tgg aca gca gcg cta ctg gag gcc ttt ggg gca gag gga cag			1219
Gly Asp Trp Thr Ala Ala Leu Leu Glu Ala Phe Gly Ala Glu Gly Gln			
360	365	370	
gcc ctc cag gag ccc tgg agc ctg cca agg ctg gca gtg gac ggg ccc			1267
Ala Leu Gln Glu Pro Trp Ser Leu Pro Arg Leu Ala Val Asp Gly Pro			
375	380	385	
ttt gga act gcc ctg aca gat gta ttt cac tac cca gtg tgt gtg tgc			1315
Phe Gly Thr Ala Leu Thr Asp Val Phe His Tyr Pro Val Cys Val Cys			
390	395	400	
gtt gcc gcg ggg atc gga gtc act ccc ttc gct gct ctt ctg aaa tct			1363
Val Ala Ala Gly Ile Gly Val Thr Pro Phe Ala Ala Leu Leu Lys Ser			



405	410	415	420	
ata tgg tac aaa tgc agt gag gca cag acc cca ctg aag ctg agc aag				1411
Ile Trp Tyr Lys Cys Ser Glu Ala Gln Thr Pro Leu Lys Leu Ser Lys				
425		430	435	
gtg tat ttc tac tgg att tgc cgg gat gca aga gct ttt gag tgg ttt				1459
Val Tyr Phe Tyr Trp Ile Cys Arg Asp Ala Arg Ala Phe Glu Trp Phe				
440	445		450	
gct gat ctc tta ctc tcc ctg gaa aca cgg atg agt gag cag ggg aaa				1507
Ala Asp Leu Leu Leu Ser Leu Glu Thr Arg Met Ser Glu Gln Gly Lys				
455	460		465	
act cac ttt ctg agt tat cat ata ttt ctt acc ggc tgg gat gaa aat				1555
Thr His Phe Leu Ser Tyr His Ile Phe Leu Thr Gly Trp Asp Glu Asn				
470	475		480	
cag gct ctt cac ata gct tta cac tgg gac gaa aat act gac gtg att				1603
Gln Ala Leu His Ile Ala Leu His Trp Asp Glu Asn Thr Asp Val Ile				
485	490	495	500	
aca ggc tta aag cag aag acc ttc tat ggg agg ccc aac tgg aac aat				1651
Thr Gly Leu Lys Gln Lys Thr Phe Tyr Gly Arg Pro Asn Trp Asn Asn				
505	510		515	
gag ttc aag cag att gcc tac aat cac ccc agc agc agt att ggc gtg				1699
Glu Phe Lys Gln Ile Ala Tyr Asn His Pro Ser Ser Ser Ile Gly Val				
520	525		530	
ttc ttc tgt gga cct aaa gct ctc tcg agg aca ctt caa aag atg tgc				1747
Phe Phe Cys Gly Pro Lys Ala Leu Ser Arg Thr Leu Gln Lys Met Cys				
535	540		545	
cac ttg tat tca tca gct gac ccc aga ggt gtt cat ttc tat tac aac				1795
His Leu Tyr Ser Ser Ala Asp Pro Arg Gly Val His Phe Tyr Tyr Asn				
550	555		560	
aag gag agc ttc tag actttggagg tcaagtccag gcattgtggt ttcaatcaag				1850
Lys Glu Ser Phe				
565				
ttattgattc caaagaactc caccaggaat tcctgtgacg gcctgttgat atgagctccc				1910
agttgggaac tgggaataa taattaacta ttgtgaacag tacactatac catacttcct				1970
tagcttataa ataacatgtc atatacaaca gaacaaaaac atttactgaa attaaaaatat				2030



attatgtttc tcca

2044

&lt;210&gt; 4

&lt;211&gt; 568

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

Met Met Gly Cys Trp Ile Leu Asn Glu Gly Leu Ser Thr Ile Leu Val  
 1 5 10 15

Leu Ser Trp Leu Gly Ile Asn Phe Tyr Leu Phe Ile Asp Thr Phe Tyr  
 20 25 30

Trp Tyr Glu Glu Glu Glu Ser Phe His Tyr Thr Arg Val Ile Leu Gly  
 35 40 45

Ser Thr Leu Ala Trp Ala Arg Ala Ser Ala Leu Cys Leu Asn Phe Asn  
 50 55 60

Cys Met Leu Ile Leu Ile Pro Val Ser Arg Asn Leu Ile Ser Phe Ile  
 65 70 75 80

Arg Gly Thr Ser Ile Cys Cys Arg Gly Pro Trp Arg Arg Gln Leu Asp  
 85 90 95

Lys Asn Leu Arg Phe His Lys Leu Val Ala Tyr Gly Ile Ala Val Asn  
 100 105 110

Ala Thr Ile His Ile Val Ala His Phe Phe Asn Leu Glu Arg Tyr His  
 115 120 125

Trp Ser Gln Ser Glu Glu Ala Gln Gly Leu Leu Ala Ala Leu Ser Lys  
 130 135 140

Leu Gly Asn Thr Pro Asn Glu Ser Tyr Leu Asn Pro Val Arg Thr Phe  
 145 150 155 160

Pro Thr Asn Thr Thr Thr Glu Leu Leu Arg Thr Ile Ala Gly Val Thr  
 165 170 175

Gly Leu Val Ile Ser Leu Ala Leu Val Leu Ile Met Thr Ser Ser Thr  
 180 185 190

Glu Phe Ile Arg Gln Ala Ser Tyr Glu Leu Phe Trp Tyr Thr His His  
 195 200 205



Val Phe Ile Val Phe Phe Leu Ser Leu Ala Ile His Gly Thr Gly Arg  
 210 215 220  
 Ile Val Arg Gly Gln Thr Gln Asp Ser Leu Ser Leu His Asn Ile Thr  
 225 230 235 240  
 Phe Cys Arg Asp Arg Tyr Ala Glu Trp Gln Thr Val Ala Gln Cys Pro  
 245 250 255  
 Val Pro Gln Phe Ser Gly Lys Glu Pro Ser Ala Trp Lys Trp Ile Leu  
 260 265 270  
 Gly Pro Val Val Leu Tyr Ala Cys Glu Arg Ile Ile Arg Phe Trp Arg  
 275 280 285  
 Phe Gln Gln Glu Val Val Ile Thr Lys Val Val Ser His Pro Ser Gly  
 290 295 300  
 Val Leu Glu Leu His Met Lys Lys Arg Gly Phe Lys Met Ala Pro Gly  
 305 310 315 320  
 Gln Tyr Ile Leu Val Gln Cys Pro Ala Ile Ser Ser Leu Glu Trp His  
 325 330 335  
 Pro Phe Thr Leu Thr Ser Ala Pro Gln Glu Asp Phe Phe Ser Val His  
 340 345 350  
 Ile Arg Ala Ala Gly Asp Trp Thr Ala Ala Leu Leu Glu Ala Phe Gly  
 355 360 365  
 Ala Glu Gly Gln Ala Leu Gln Glu Pro Trp Ser Leu Pro Arg Leu Ala  
 370 375 380  
 Val Asp Gly Pro Phe Gly Thr Ala Leu Thr Asp Val Phe His Tyr Pro  
 385 390 395 400  
 Val Cys Val Cys Val Ala Ala Gly Ile Gly Val Thr Pro Phe Ala Ala  
 405 410 415  
 Leu Leu Lys Ser Ile Trp Tyr Lys Cys Ser Glu Ala Gln Thr Pro Leu  
 420 425 430  
 Lys Leu Ser Lys Val Tyr Phe Tyr Trp Ile Cys Arg Asp Ala Arg Ala  
 435 440 445  
 Phe Glu Trp Phe Ala Asp Leu Leu Leu Ser Leu Glu Thr Arg Met Ser  
 450 455 460



Glu Gln Gly Lys Thr His Phe Leu Ser Tyr His Ile Phe Leu Thr Gly  
 465 470 475 480

Trp Asp Glu Asn Gln Ala Leu His Ile Ala Leu His Trp Asp Glu Asn  
 485 490 495

Thr Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Phe Tyr Gly Arg Pro  
 500 505 510

Asn Trp Asn Asn Glu Phe Lys Gln Ile Ala Tyr Asn His Pro Ser Ser  
 515 520 525

Ser Ile Gly Val Phe Phe Cys Gly Pro Lys Ala Leu Ser Arg Thr Leu  
 530 535 540

Gln Lys Met Cys His Leu Tyr Ser Ser Ala Asp Pro Arg Gly Val His  
 545 550 555 560

Phe Tyr Tyr Asn Lys Glu Ser Phe  
 565

<210> 5  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer

<400> 5  
 aacaagcgtg gcttcagcat g 21

<210> 6  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer

<400> 6  
 agcaatattg ttggtcat 18

<210> 7  
 <211> 24



<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 7

gacttgacag aaaatctata aggg

24

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 8

ttgtaccaga tggatttcaa

20

<210> 9

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 9

caggtctgaa acagaaaacc t

21

<210> 10

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 10

atgaattctc attaattatt caataaa

27

<210> 11

<211> 20



&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 11

atctcaaaag actctgcaca

20

&lt;210&gt; 12

&lt;211&gt; 569

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

Gly Asn Trp Ala Val Asn Glu Gly Leu Ser Ile Phe Ala Ile Leu Val  
 1 5 10 15

Trp Leu Gly Leu Asn Val Phe Leu Phe Val Trp Tyr Tyr Arg Val Tyr  
 20 25 30

Asp Ile Pro Pro Lys Phe Phe Tyr Thr Arg Lys Leu Leu Gly Ser Ala  
 35 40 45

Leu Ala Leu Ala Arg Ala Pro Ala Ala Cys Leu Asn Phe Asn Cys Met  
 50 55 60

Leu Ile Leu Leu Pro Val Cys Arg Asn Leu Leu Ser Phe Leu Arg Gly  
 65 70 75 80

Ser Ser Ala Cys Cys Ser Thr Arg Val Arg Arg Gln Leu Asp Arg Asn  
 85 90 95

Leu Thr Phe His Lys Met Val Ala Trp Met Ile Ala Leu His Ser Ala  
 100 105 110

Ile His Thr Ile Ala His Leu Phe Asn Val Glu Trp Cys Val Asn Ala  
 115 120 125

Arg Val Asn Asn Ser Asp Pro Tyr Ser Val Ala Leu Ser Glu Leu Gly  
 130 135 140

Asp Arg Gln Asn Glu Ser Tyr Leu Asn Phe Ala Arg Lys Arg Ile Lys  
 145 150 155 160

Asn Pro Glu Gly Gly Leu Tyr Leu Ala Val Thr Leu Leu Ala Gly Ile  
 165 170 175



Thr Gly Val Val Ile Thr Leu Cys Leu Ile Leu Ile Ile Thr Ser Ser  
 180 185 190  
 Thr Lys Thr Ile Arg Arg Ser Tyr Phe Glu Val Phe Trp Tyr Thr His  
 195 200 205  
 His Leu Phe Val Ile Phe Phe Ile Gly Leu Ala Ile His Gly Ala Glu  
 210 215 220  
 Arg Ile Val Arg Gly Gln Thr Ala Glu Ser Leu Ala Val His Asn Ile  
 225 230 235 240  
 Thr Val Cys Glu Gln Lys Ile Ser Glu Trp Gly Lys Ile Lys Glu Cys  
 245 250 255  
 Pro Ile Pro Gln Phe Ala Gly Asn Pro Pro Met Thr Trp Lys Trp Ile  
 260 265 270  
 Val Gly Pro Met Phe Leu Tyr Leu Cys Glu Arg Leu Val Arg Phe Trp  
 275 280 285  
 Arg Ser Gln Gln Lys Val Val Ile Thr Lys Val Val Thr His Pro Phe  
 290 295 300  
 Lys Thr Ile Glu Leu Gln Met Lys Lys Lys Gly Phe Lys Met Glu Val  
 305 310 315 320  
 Gly Gln Tyr Ile Phe Val Lys Cys Pro Lys Val Ser Lys Leu Glu Trp  
 325 330 335  
 His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Asp Phe Phe Ser Ile  
 340 345 350  
 His Ile Arg Ile Val Gly Asp Trp Thr Glu Gly Leu Phe Asn Ala Cys  
 355 360 365  
 Gly Cys Asp Lys Gln Glu Phe Gln Asp Ala Trp Lys Leu Pro Lys Ile  
 370 375 380  
 Ala Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val Phe Ser Tyr  
 385 390 395 400  
 Glu Val Val Met Leu Val Gly Ala Gly Ile Gly Val Thr Pro Phe Ala  
 405 410 415  
 Ser Ile Leu Lys Ser Val Trp Tyr Lys Tyr Cys Asn Asn Ala Thr Asn  
 420 425 430



Leu Lys Leu Lys Lys Ile Tyr Phe Tyr Trp Leu Cys Arg Asp Thr His  
 435 440 445

Ala Phe Glu Trp Phe Ala Asp Leu Leu Gln Leu Leu Glu Ser Gln Met  
 450 455 460

Gln Glu Arg Asn Asn Ala Gly Phe Leu Ser Tyr Asn Ile Tyr Leu Thr  
 465 470 475 480

Gly Trp Asp Glu Ser Gln Ala Asn His Phe Ala Val His His Asp Glu  
 485 490 495

Glu Lys Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Leu Tyr Gly Arg  
 500 505 510

Pro Asn Trp Asp Asn Glu Phe Lys Thr Ile Ala Ser Gln His Pro Asn  
 515 520 525

Thr Arg Ile Gly Val Phe Leu Cys Gly Pro Glu Ala Leu Ala Glu Thr  
 530 535 540

Leu Ser Lys Gln Ser Ile Ser Asn Ser Glu Ser Gly Pro Arg Gly Val  
 545 550 555 560

His Phe Ile Phe Asn Lys Glu Asn Phe  
 565

<210> 13

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 13

ttggctaaat cccatcca

18

<210> 14

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer



&lt;400&gt; 14

tgcatgacca acaatattgc t

21

&lt;210&gt; 15

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 15

caaggtacct cttgaccatg ggaaact

27

&lt;210&gt; 16

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 16

acgaattcaa gtaaattact gaagata

27

&lt;210&gt; 17

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; modified\_base

&lt;222&gt; (())

&lt;223&gt; n at position 3 = inosine

&lt;220&gt;

&lt;221&gt; modified\_base

&lt;222&gt; (())

&lt;223&gt; n at position 6 = inosine

&lt;220&gt;

&lt;221&gt; modified\_base

&lt;222&gt; (())

&lt;223&gt; n at position 12 = inosine



<220>

<223> Description of Artificial Sequence: primer

<400> 17

ccngtntgtc gnaatctgct stcctt

26

<210> 18

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<221> modified\_base

<222> (5)

<223> n at position 5 = inosine

<220>

<221> modified\_base

<222> (9)

<223> n at position 9 = inosine

<220>

<221> modified\_base

<222> (11)

<223> n at position 11 = inosine

<220>

<223> Description of Artificial Sequence: primer

<400> 18

tcccngcana nccagtagaa rtagatctt

29

<210> 19

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 19

ttggcacagt cagtgaggat gtcttc

26

<210> 20



&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 20

ctgttggtt ctactgtac gttcaaagtt

30

&lt;210&gt; 21

&lt;211&gt; 563

&lt;212&gt; PRT

&lt;213&gt; Rat

&lt;400&gt; 21

Met	Gly	Asn	Trp	Leu	Val	Asn	His	Trp	Leu	Ser	Val	Leu	Phe	Leu	Val
1				5					10					15	

Ser	Trp	Leu	Gly	Leu	Asn	Ile	Phe	Leu	Phe	Val	Tyr	Val	Phe	Leu	Asn
		20						25					30		

Tyr	Glu	Lys	Ser	Asp	Lys	Tyr	Tyr	Tyr	Thr	Arg	Glu	Ile	Leu	Gly	Thr
		35					40					45			

Ala	Leu	Ala	Leu	Ala	Arg	Ala	Ser	Ala	Leu	Cys	Leu	Asn	Phe	Asn	Ser
	50					55					60				

Met	Val	Ile	Leu	Ile	Pro	Val	Cys	Arg	Asn	Leu	Leu	Ser	Phe	Leu	Arg
65					70					75					80

Gly	Thr	Cys	Ser	Phe	Cys	Asn	His	Thr	Leu	Arg	Lys	Pro	Leu	Asp	His
			85						90					95	

Asn	Leu	Thr	Phe	His	Lys	Leu	Val	Ala	Tyr	Met	Ile	Cys	Ile	Phe	Thr
		100						105					110		

Ala	Ile	His	Ile	Ile	Ala	His	Leu	Phe	Asn	Phe	Glu	Arg	Tyr	Ser	Arg
	115						120				125				

Ser	Gln	Gln	Ala	Met	Asp	Gly	Ser	Leu	Ala	Ser	Val	Leu	Ser	Ser	Leu
	130					135					140				

Phe	His	Pro	Glu	Lys	Glu	Asp	Ser	Trp	Leu	Asn	Pro	Ile	Gln	Ser	Pro
145				150					155					160	

Asn	Val	Thr	Val	Met	Tyr	Ala	Ala	Phe	Thr	Ser	Ile	Ala	Gly	Leu	Thr
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----



	165		170		175
Gly Val Val Ala Thr Val Ala Leu Val Leu Met Val Thr Ser Ala Met					
180		185		190	
Glu Phe Ile Arg Arg Asn Tyr Phe Glu Leu Phe Trp Tyr Thr His His					
195		200		205	
Leu Phe Ile Ile Tyr Ile Ile Cys Leu Gly Ile His Gly Leu Gly Gly					
210		215		220	
Ile Val Arg Gly Gln Thr Glu Glu Ser Met Ser Glu Ser His Pro Arg					
225		230		235	240
Asn Cys Ser Tyr Ser Phe His Glu Trp Asp Lys Tyr Glu Arg Ser Cys					
	245		250		255
Arg Ser Pro His Phe Val Gly Gln Pro Pro Glu Ser Trp Lys Trp Ile					
	260		265		270
Leu Ala Pro Ile Ala Phe Tyr Ile Phe Glu Arg Ile Leu Arg Phe Tyr					
	275		280		285
Arg Ser Arg Gln Lys Val Val Ile Thr Lys Val Val Met His Pro Cys					
	290		295		300
Lys Val Leu Glu Leu Gln Met Arg Lys Arg Gly Phe Thr Met Gly Ile					
305		310		315	320
Gly Gln Tyr Ile Phe Val Asn Cys Pro Ser Ile Ser Phe Leu Glu Trp					
	325		330		335
His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Glu Phe Phe Ser Ile					
	340		345		350
His Ile Arg Ala Ala Gly Asp Trp Thr Glu Asn Leu Ile Arg Thr Phe					
	355		360		365
Glu Gln Gln His Ser Pro Met Pro Arg Ile Glu Val Asp Gly Pro Phe					
	370		375		380
Gly Thr Val Ser Glu Asp Val Phe Gln Tyr Glu Val Ala Val Leu Val					
385		390		395	400
Gly Ala Gly Ile Gly Val Thr Pro Phe Ala Ser Phe Leu Lys Ser Ile					
	405		410		415
Trp Tyr Lys Phe Gln Arg Ala His Asn Lys Leu Lys Thr Gln Lys Ile					



420                      425                      430  
 Tyr Phe Tyr Trp Ile Cys Arg Glu Thr Gly Ala Phe Ala Trp Phe Asn  
     435                      440                      445  
 Asn Leu Leu Asn Ser Leu Glu Gln Glu Met Asp Glu Leu Gly Lys Pro  
     450                      455                      460  
 Asp Phe Leu Asn Tyr Arg Leu Phe Leu Thr Gly Trp Asp Ser Asn Ile  
     465                      470                      475                      480  
 Ala Gly His Ala Ala Leu Asn Phe Asp Arg Ala Thr Asp Val Leu Thr  
                     485                      490                      495  
 Gly Leu Lys Gln Lys Thr Ser Phe Gly Arg Pro Met Trp Asp Asn Glu  
                     500                      505                      510  
 Phe Ser Arg Ile Ala Thr Ala His Pro Lys Ser Val Val Gly Val Phe  
                     515                      520                      525  
 Leu Cys Gly Pro Pro Thr Leu Ala Lys Ser Leu Arg Lys Cys Cys Arg  
                     530                      535                      540  
 Arg Tyr Ser Ser Leu Asp Pro Arg Lys Val Gln Phe Tyr Phe Asn Lys  
     545                      550                      555                      560  
 Glu Thr Phe

<210> 22  
 <211> 2577  
 <212> DNA  
 <213> Rat

<400> 22  
 ttctgagtag gtgtgcattt gagggtcata aagacatata tcttgagcta gacagaagtt 60  
 cctatcctga aggatcccat cagagaaacc agattgctcc taagaggctc cagacctcca 120  
 tttgacaatg ggaaactggc tggtaacca ctggctctca gttttgtttc tggtttcttg 180  
 gttggggctg aacatttttc tgtttgtgta cgtcttcctg aattatgaga agtctgacaa 240  
 gtactattac acgagagaaa ttctcggaac tgccttggcc ttggccagag catctgcttt 300  
 gtgcctgaat tttacagca tggatgcct gattcctgtg tgcgaaatc tgctctcctt 360



cctgaggggc acctgctcat ttgcaacca cacgctgaga aagccattgg atcacaacct 420  
caccttccat aagctgggtg catatatgat ctgcatattc acagctattc atatcattgc 480  
acatctattt aactttgaac gctacagtag aagccaacag gccatggatg gatctcttgc 540  
ctctgttctc tccagcctat tccatcccga gaaagaagat tcttggtctaa atcccatcca 600  
gtctccaaac gtgacagtga tgtatgcagc atttaccagt attgctggcc ttactggagt 660  
ggtcgccact gtggcttttg ttctcatggt aacttcagct atggagttaa tccgcaggaa 720  
ttattttgag ctcttctggt atacacatca ctttttcac atctatatca tctgcttagg 780  
gatccatggc ctggggggga ttgtccgggg tcaaacagaa gagagcatga gtgaaagtca 840  
tccccgcaac tggtcatact ctttccacga gtgggataag tatgaaagga gttgcaggag 900  
tcctcatttt gtggggcaac ccctgagtc ttggaagtgg atcctcgccg cgattgcttt 960  
ttatatcttt gaaaggatcc ttcgctttta tcgctcccg cagaaggctg tgattaccaa 1020  
ggttgatcag caccatgta aagttttgga attgcagatg aggaagcggg gctttactat 1080  
gggaatagga cagtatatat tcgtaaattg ccctcgatt tecttcctgg aatggcatcc 1140  
ctttactctg acctctgctc cagaggaaga attttctcc attcatattc gagcagcagg 1200  
ggactggaca gaaaatctca taaggacatt tgaacaacag cactcaccaa tgcccaggat 1260  
cgaggtggat ggtccctttg gcacagtcag tgaggatgct ttccagtacg aagtggctgt 1320  
actggttggg gcagggattg gcgtcactcc ctttgcttcc ttcttgaaat ctatctggta 1380  
caaattccag cgtgcacaca acaagctgaa aacacaaaag atctatttct actggatttg 1440  
tagagagacg ggtgcctttg cctggttcaa caacttattg aattccctgg aacaagagat 1500  
ggacgaatta ggcaaaccgg atttcctaaa ctaccgactc ttcctcactg gctgggatag 1560  
caacattgct ggtcatgcag cattaaactt tgacagagcc actgacgtcc tgacaggtct 1620  
gaaacagaaa acctcctttg ggagaccaat gtgggacaat gagttttcta gaatagctac 1680  
tgcccacccc aagtctgtgg tgggggtttt cttatgcggc cctccgactt tggcaaaaag 1740  
cctgcgcaaa tgctgtcggc ggtactcaag tctggatcct aggaagggtc aattctactt 1800



caacaaagaa acgttctgaa ttggaggaag cgcacagta gtacttctcc atcttccttt 1860  
 tcactaacgt gtgggtcagc tactagatag tccgttgctg cacaaggact tcactcccat 1920  
 cttaaagttg actcaactcc atcattcttg ggctttggca acatgagagc tgcataactc 1980  
 acaattgcaa aacacatgaa ttattattgg ggggattgta aatccttctg ggaaacctgc 2040  
 ctttagctga atcttgctgg ttgacacttg cacaatttaa cctcaggtgt cttggttgat 2100  
 acctgataat cttccctccc acctgtccct cacagaagat ttctaagtag ggtgatttta 2160  
 aaatatttat tgaatccacg acaaaacaat aatcataaat aataaacata aaattaccaa 2220  
 gattcccact cccatatcat acccactaag aacatcgta tacatgagct tatcatccag 2280  
 tgtgaccaac aatttatact ttactgtgcc aaaataatct tcatctttgc ttattgaaca 2340  
 attttgctga ctttccttag taatatctta agtatattaa ctggaatcaa atttgtatta 2400  
 tagttagaag ccaactatat tgccagtttg tattgtttga aataactgga aaggcctgac 2460  
 ctacatcggtg gggtaattta acagaagctc tttccatttt ttgttggtgt tgttaaagag 2520  
 ttttgtttat gaatgtgtta taaaaagaaa ataaaaagtt ataattttga .cggaaaa 2577

&lt;210&gt; 23

&lt;211&gt; 332

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

Glu	Ser	Met	Asn	Glu	Ser	His	Pro	Arg	Lys	Cys	Ala	Glu	Ser	Phe	Glu
1				5					10					15	
Met	Trp	Asp	Asp	Arg	Asp	Ser	His	Cys	Arg	Arg	Pro	Lys	Phe	Glu	Gly
		20						25					30		
His	Pro	Pro	Glu	Ser	Trp	Lys	Trp	Ile	Leu	Ala	Pro	Val	Ile	Leu	Tyr
		35					40					45			
Ile	Cys	Glu	Arg	Ile	Leu	Arg	Phe	Tyr	Arg	Ser	Gln	Gln	Lys	Val	Val
	50					55					60				
Ile	Thr	Lys	Val	Val	Met	His	Pro	Ser	Lys	Val	Leu	Glu	Leu	Gln	Met
65					70					75				80	



Asn Lys Arg Gly Phe Ser Met Glu Val Gly Gln Tyr Ile Phe Val Asn  
                             85                            90                            95  
 Cys Pro Ser Ile Ser Leu Leu Glu Trp His Pro Phe Thr Leu Thr Ser  
                             100                            105                            110  
 Ala Pro Glu Glu Asp Phe Phe Ser Ile His Ile Arg Ala Ala Gly Asp  
                             115                            120                            125  
 Trp Thr Glu Asn Leu Ile Arg Ala Phe Glu Gln Gln Tyr Ser Pro Ile  
                             130                            135                            140  
 Pro Arg Ile Glu Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val  
                             145                            150                            155                            160  
 Phe Gln Tyr Glu Val Ala Val Leu Val Gly Ala Gly Ile Gly Val Thr  
                             165                            170                            175  
 Pro Phe Ala Ser Ile Leu Lys Ser Ile Trp Tyr Lys Phe Gln Cys Ala  
                             180                            185                            190  
 Asp His Asn Leu Lys Thr Lys Lys Ile Tyr Phe Tyr Trp Ile Cys Arg  
                             195                            200                            205  
 Glu Thr Gly Ala Phe Ser Trp Phe Asn Asn Leu Leu Thr Ser Leu Glu  
                             210                            215                            220  
 Gln Glu Met Glu Glu Leu Gly Lys Val Gly Phe Leu Asn Tyr Arg Leu  
                             225                            230                            235                            240  
 Phe Leu Thr Gly Trp Asp Ser Asn Ile Val Gly His Ala Ala Leu Asn  
                             245                            250                            255  
 Phe Asp Lys Ala Thr Asp Ile Val Thr Gly Leu Lys Gln Lys Thr Ser  
                             260                            265                            270  
 Phe Gly Arg Pro Met Trp Asp Asn Glu Phe Ser Thr Ile Ala Thr Ser  
                             275                            280                            285  
 His Pro Lys Ser Val Val Gly Val Phe Leu Cys Gly Pro Arg Thr Leu  
                             290                            295                            300  
 Ala Lys Ser Leu Arg Lys Cys Cys His Arg Tyr Ser Ser Leu Asp Pro  
                             305                            310                            315                            320  
 Arg Lys Val Gln Phe Tyr Phe Asn Lys Glu Asn Phe  
                             325                            330



<210> 24  
<211> 14  
<212> PRT  
<213> Homo sapiens

<400> 24  
Cys Ala Glu Ser Phe Glu Met Trp Asp Asp Arg Asp Ser His  
1 5 10

<210> 25  
<211> 14  
<212> PRT  
<213> Homo sapiens

<400> 25  
Lys Ser Leu Arg Lys Cys Cys His Arg Tyr Ser Ser Leu Asp  
1 5 10

<210> 26  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer

<400> 26  
gaagtgggtgg gaggcgaaga cata 24

<210> 27  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer

<400> 27  
cctgtcatatc ctgggacggt ctgg 24

<210> 28  
<211> 24  
<212> DNA



<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 28

gagcacagtg agatgcctgt tcag

24

<210> 29

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 29

ggaaggcagc agagagcaat gatg

24

<210> 30

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 30

acatctgcga gcggcacttc caga

24

<210> 31

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 31

agctcgtcaa caggcaggac cgagc

25

<210> 32

<211> 24

<212> DNA



<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 32

gcagtgcac cacaatcttca gcac

24

<210> 33

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 33

gagagctctg gagacacttg agttc

25

<210> 34

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 34

catgttctct ctggctgaca ag

22

<210> 35

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 35

cacaatagcg agctccgctt cacgc

25

<210> 36

<211> 24

<212> DNA



<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 36

gcaggacatc aaccctgcac tctc

24

<210> 37

<211> 570

<212> PRT

<213> Bovine

<400> 37

Met Gly Asn Trp Val Val Asn Glu Gly Ile Ser Ile Phe Val Ile Leu  
1 5 10 15

Val Trp Leu Gly Met Asn Val Phe Leu Phe Val Trp Tyr Tyr Arg Val  
20 25 30

Tyr Asp Ile Pro Asp Lys Phe Phe Tyr Thr Arg Lys Leu Leu Gly Ser  
35 40 45

Ala Leu Ala Leu Ala Arg Ala Pro Ala Ala Cys Leu Asn Phe Asn Cys  
50 55 60

Met Leu Ile Leu Leu Pro Val Cys Arg Asn Leu Leu Ser Phe Leu Arg  
65 70 75 80

Gly Ser Ser Ala Cys Cys Ser Thr Arg Ile Arg Arg Gln Leu Asp Arg  
85 90 95

Asn Leu Thr Phe His Lys Met Val Ala Trp Met Ile Ala Leu His Thr  
100 105 110

Ala Ile His Thr Ile Ala His Leu Phe Asn Val Glu Trp Cys Val Asn  
115 120 125

Ala Arg Val Asn Asn Ser Asp Pro Tyr Ser Ile Ala Leu Ser Asp Ile  
130 135 140

Gly Asp Lys Pro Asn Glu Thr Tyr Leu Asn Phe Val Arg Gln Arg Ile  
145 150 155 160

Lys Asn Pro Glu Gly Gly Leu Tyr Val Ala Val Thr Arg Leu Ala Gly  
165 170 175



Ile Thr Gly Val Val Ile Thr Leu Cys Leu Ile Leu Ile Ile Thr Ser  
 180 185 190  
 Ser Thr Lys Thr Ile Arg Arg Ser Tyr Phe Glu Val Phe Trp Tyr Thr  
 195 200 205  
 His His Leu Phe Val Ile Phe Phe Ile Gly Leu Ala Ile His Gly Ala  
 210 215 220  
 Gln Arg Ile Val Arg Gly Gln Thr Ala Glu Ser Leu Leu Lys His Gln  
 225 230 235 240  
 Pro Arg Asn Cys Tyr Gln Asn Ile Ser Gln Trp Gly Lys Ile Glu Asn  
 245 250 255  
 Cys Pro Ile Pro Glu Phe Ser Gly Asn Pro Pro Met Thr Trp Lys Trp  
 260 265 270  
 Ile Val Gly Pro Met Phe Leu Tyr Leu Cys Glu Arg Leu Val Arg Phe  
 275 280 285  
 Trp Arg Ser Gln Gln Lys Val Val Ile Thr Lys Val Val Thr His Pro  
 290 295 300  
 Phe Lys Thr Ile Glu Leu Gln Met Lys Lys Lys Gly Phe Lys Met Glu  
 305 310 315 320  
 Val Gly Gln Tyr Ile Phe Val Lys Cys Pro Val Val Ser Lys Leu Glu  
 325 330 335  
 Trp His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Asp Phe Phe Ser  
 340 345 350  
 Ile His Ile Arg Ile Val Gly Asp Trp Thr Glu Gly Leu Phe Lys Ala  
 355 360 365  
 Cys Gly Cys Asp Lys Gln Glu Phe Gln Asp Ala Trp Lys Leu Pro Lys  
 370 375 380  
 Ile Ala Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val Phe Ser  
 385 390 395 400  
 Tyr Glu Val Val Met Leu Val Gly Ala Gly Ile Gly Val Thr Pro Phe  
 405 410 415  
 Ala Ser Ile Leu Lys Ser Val Trp Tyr Lys Tyr Cys Asn Lys Ala Pro  
 420 425 430



Asn Leu Arg Leu Lys Lys Ile Tyr Phe Tyr Trp Leu Cys Arg Asp Thr  
 435 440 445

His Ala Phe Glu Trp Phe Ala Asp Leu Leu Gln Leu Leu Glu Thr Gln  
 450 455 460

Met Gln Glu Lys Asn Asn Thr Asp Phe Leu Ser Tyr Asn Ile Cys Leu  
 465 470 475 480

Thr Gly Trp Asp Glu Ser Gln Ala Ser His Phe Ala Met His His Asp  
 485 490 495

Glu Glu Lys Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Leu Tyr Gly  
 500 505 510

Arg Pro Asn Trp Asp Asn Glu Phe Lys Thr Ile Gly Ser Gln His Pro  
 515 520 525

Asn Thr Arg Ile Gly Val Phe Leu Cys Gly Pro Glu Ala Leu Ala Asp  
 530 535 540

Thr Leu Asn Lys Gln Cys Ile Ser Asn Ser Asp Ser Gly Pro Arg Gly  
 545 550 555 560

Val His Phe Ile Phe Asn Lys Glu Asn Phe  
 565 570

<210> 38

<211> 570

<212> PRT

<213> murine

<400> 38

Met Gly Asn Trp Ala Val Asn Glu Gly Leu Ser Ile Phe Val Ile Leu  
 1 5 10 15

Val Trp Leu Gly Leu Asn Val Phe Leu Phe Ile Asn Tyr Tyr Lys Val  
 20 25 30

Tyr Asp Asp Gly Pro Lys Tyr Asn Tyr Thr Arg Lys Leu Leu Gly Ser  
 35 40 45

Ala Leu Ala Leu Ala Arg Ala Pro Ala Ala Cys Leu Asn Phe Asn Cys  
 50 55 60

Met Leu Ile Leu Leu Pro Val Cys Arg Asn Leu Leu Ser Phe Leu Arg  
 65 70 75 80



Gly Ser Ser Ala Cys Cys Ser Thr Arg Ile Arg Arg Gln Leu Asp Arg  
                     85                    90                    95  
 Asn Leu Thr Phe His Lys Met Val Ala Trp Met Ile Ala Leu His Thr  
                     100                    105                    110  
 Ala Ile His Thr Ile Ala His Leu Phe Asn Val Glu Trp Cys Val Asn  
                     115                    120                    125  
 Ala Arg Val Gly Ile Ser Asp Arg Tyr Ser Ile Ala Leu Ser Asp Ile  
                     130                    135                    140  
 Gly Asp Asn Glu Asn Glu Glu Tyr Leu Asn Phe Ala Arg Glu Lys Ile  
 145                    150                    155                    160  
 Lys Asn Pro Glu Gly Gly Leu Tyr Val Ala Val Thr Arg Leu Ala Gly  
                     165                    170                    175  
 Ile Thr Gly Ile Val Ile Thr Leu Cys Leu Ile Leu Ile Ile Thr Ser  
                     180                    185                    190  
 Ser Thr Lys Thr Ile Arg Arg Ser Tyr Phe Glu Val Phe Trp Tyr Thr  
                     195                    200                    205  
 His His Leu Phe Val Ile Phe Phe Ile Gly Leu Ala Ile His Gly Ala  
                     210                    215                    220  
 Glu Arg Ile Val Arg Gly Gln Thr Ala Glu Ser Leu Glu Glu His Asn  
 225                    230                    235                    240  
 Leu Asp Ile Cys Ala Asp Lys Ile Glu Glu Trp Gly Lys Ile Lys Glu  
                     245                    250                    255  
 Cys Pro Val Pro Lys Phe Ala Gly Asn Pro Pro Met Thr Trp Lys Trp  
                     260                    265                    270  
 Ile Val Gly Pro Met Phe Leu Tyr Leu Cys Glu Arg Leu Val Arg Phe  
                     275                    280                    285  
 Trp Arg Ser Gln Gln Lys Val Val Ile Thr Lys Val Val Thr His Pro  
                     290                    295                    300  
 Phe Lys Thr Ile Glu Leu Gln Met Lys Lys Lys Gly Phe Lys Met Glu  
 305                    310                    315                    320  
 Val Gly Gln Tyr Ile Phe Val Lys Cys Pro Lys Val Ser Lys Leu Glu  
                     325                    330                    335



Trp His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Asp Phe Phe Ser  
 340 345 350  
 Ile His Ile Arg Ile Val Gly Asp Trp Thr Glu Gly Leu Phe Asn Ala  
 355 360 365  
 Cys Gly Cys Asp Lys Gln Glu Phe Gln Asp Ala Trp Lys Leu Pro Lys  
 370 375 380  
 Ile Ala Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val Phe Ser  
 385 390 395 400  
 Tyr Glu Val Val Met Leu Val Gly Ala Gly Ile Gly Val Thr Pro Phe  
 405 410 415  
 Ala Ser Ile Leu Lys Ser Val Trp Tyr Lys Tyr Cys Asp Asn Ala Thr  
 420 425 430  
 Ser Leu Lys Leu Lys Lys Ile Tyr Phe Tyr Trp Leu Cys Arg Asp Thr  
 435 440 445  
 His Ala Phe Glu Trp Phe Ala Asp Leu Leu Gln Leu Leu Glu Thr Gln  
 450 455 460  
 Met Gln Glu Arg Asn Asn Ala Asn Phe Leu Ser Tyr Asn Ile Tyr Leu  
 465 470 475 480  
 Thr Gly Trp Asp Glu Ser Gln Ala Asn His Phe Ala Val His His Asp  
 485 490 495  
 Glu Glu Lys Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Leu Tyr Gly  
 500 505 510  
 Arg Pro Asn Trp Asp Asn Glu Phe Lys Thr Ile Ala Ser Glu His Pro  
 515 520 525  
 Asn Thr Thr Ile Gly Val Phe Leu Cys Gly Pro Glu Ala Leu Ala Glu  
 530 535 540  
 Thr Leu Ser Lys Gln Ser Ile Ser Asn Ser Glu Ser Gly Pro Arg Gly  
 545 550 555 560  
 Val His Phe Ile Phe Asn Lys Glu Asn Phe  
 565 570

&lt;210&gt; 39



&lt;211&gt; 944

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis sp.

&lt;400&gt; 39

```

Met Lys Pro Phe Ser Lys Asn Asp Arg Arg Arg Trp Ser Phe Asp Ser
 1             5             10             15

Val Ser Ala Gly Lys Thr Ala Val Gly Ser Ala Ser Thr Ser Pro Gly
          20             25             30

Thr Glu Tyr Ser Ile Asn Gly Asp Gln Glu Phe Val Glu Val Thr Ile
 35             40             45

Asp Leu Gln Asp Asp Asp Thr Ile Val Leu Arg Ser Val Glu Pro Ala
 50             55             60

Thr Ala Ile Asn Val Ile Gly Asp Ile Ser Asp Asp Asn Thr Gly Ile
 65             70             75             80

Met Thr Pro Val Ser Ile Ser Arg Ser Pro Thr Met Lys Arg Thr Ser
          85             90             95

Ser Asn Arg Phe Arg Gln Phe Ser Gln Glu Leu Lys Ala Glu Ala Val
          100             105             110

Ala Lys Ala Lys Gln Leu Ser Gln Glu Leu Lys Arg Phe Ser Trp Ser
          115             120             125

Arg Ser Phe Ser Gly Asn Leu Thr Thr Thr Ser Thr Ala Ala Asn Gln
          130             135             140

Ser Gly Gly Ala Gly Gly Gly Leu Val Asn Ser Ala Leu Glu Ala Arg
          145             150             155             160

Ala Leu Arg Lys Gln Arg Ala Gln Leu Asp Arg Thr Arg Ser Ser Ala
          165             170             175

Gln Arg Ala Leu Arg Gly Leu Arg Phe Ile Ser Asn Lys Gln Lys Asn
          180             185             190

Val Asp Gly Trp Asn Asp Val Gln Ser Asn Phe Glu Lys Phe Glu Lys
          195             200             205

Asn Gly Tyr Ile Tyr Arg Ser Asp Phe Ala Gln Cys Ile Gly Met Lys
          210             215             220

Asp Ser Lys Glu Phe Ala Leu Glu Leu Phe Asp Ala Leu Ser Arg Arg

```



225	230	235	240
Arg Arg Leu Lys Val Glu Lys Ile Asn His Asp Glu Leu Tyr Glu Tyr	245	250	255
Trp Ser Gln Ile Asn Asp Glu Ser Phe Asp Ser Arg Leu Gln Ile Phe	260	265	270
Phe Asp Ile Val Asp Lys Asn Glu Asp Gly Arg Ile Thr Glu Glu Glu	275	280	285
Val Lys Glu Ile Ile Met Leu Ser Ala Ser Ala Asn Lys Leu Ser Arg	290	295	300
Leu Lys Glu Gln Ala Glu Glu Tyr Ala Ala Leu Ile Met Glu Glu Leu	305	310	315
Asp Pro Glu Arg Leu Gly Tyr Ile Glu Leu Trp Gln Leu Glu Thr Leu	325	330	335
Leu Leu Gln Lys Asp Thr Tyr Leu Asn Tyr Ser Gln Ala Leu Ser Tyr	340	345	350
Thr Ser Gln Ala Leu Ser Gln Asn Leu Gln Gly Leu Arg Gly Lys Ser	355	360	365
Arg Ile His Arg Met Ser Ser Asp Phe Val Tyr Ile Met Gln Glu Asn	370	375	380
Trp Lys Arg Ile Trp Val Leu Ser Leu Trp Ile Met Ile Met Ile Gly	385	390	395
Leu Phe Leu Trp Lys Phe Phe Gln Tyr Lys Gln Lys Asp Ala Phe His	405	410	415
Val Met Gly Tyr Cys Leu Leu Thr Ala Lys Gly Ala Ala Glu Thr Leu	420	425	430
Lys Phe Asn Met Ala Leu Ile Leu Phe Pro Val Cys Arg Asn Thr Ile	435	440	445
Thr Trp Leu Arg Ser Thr Arg Leu Ser Tyr Phe Val Pro Phe Asp Asp	450	455	460
Asn Ile Asn Phe His Lys Thr Ile Ala Gly Ala Ile Val Val Ala Val	465	470	475
Ile Leu His Ile Gly Asp His Leu Ala Cys Asp Phe Pro Arg Ile Val			



36



740	745	750
Gly Leu Gly Ile Gly Ala Thr Pro Phe Ile Ser Ile Leu Lys Asp Leu		
755	760	765
Leu Asn Asn Ile Val Lys Met Glu Glu His Ala Asp Ser Ile Ser Asp		
770	775	780
Phe Ser Arg Ser Ser Glu Tyr Ser Thr Gly Ser Asn Gly Asp Thr Pro		
785	790	795
Arg Arg Lys Arg Ile Leu Lys Thr Thr Asn Ala Tyr Phe Tyr Trp Val		
805	810	815
Thr Arg Glu Gln Gly Ser Phe Asp Trp Phe Lys Gly Val Met Asn Glu		
820	825	830
Val Ala Glu Leu Asp Gln Arg Gly Val Ile Glu Met His Asn Tyr Leu		
835	840	845
Thr Ser Val Tyr Glu Glu Gly Asp Ala Arg Ser Ala Leu Ile Thr Met		
850	855	860
Val Gln Ala Leu Asn His Ala Lys Asn Gly Val Asp Ile Val Ser Gly		
865	870	875
Thr Arg Val Arg Thr His Phe Ala Arg Pro Asn Trp Lys Lys Val Leu		
885	890	895
Thr Lys Leu Ser Ser Lys His Cys Asn Ala Arg Ile Gly Val Phe Tyr		
900	905	910
Cys Gly Val Pro Val Leu Gly Lys Glu Leu Ser Lys Leu Cys Asn Thr		
915	920	925
Phe Asn Gln Lys Gly Ser Thr Lys Phe Glu Phe His Lys Glu His Phe		
930	935	940

&lt;210&gt; 40

&lt;211&gt; 590

&lt;212&gt; PRT

&lt;213&gt; Rice

&lt;400&gt; 40



Asn Leu Ala Gly Leu Arg Lys Lys Ser Ser Ile Arg Lys Ile Ser Thr  
 1 5 10 15  
 Ser Leu Ser Tyr Tyr Phe Glu Asp Asn Trp Lys Arg Leu Trp Val Leu  
 20 25 30  
 Ala Leu Trp Ile Gly Ile Met Ala Gly Leu Phe Thr Trp Lys Phe Met  
 35 40 45  
 Gln Tyr Arg Asn Arg Tyr Val Phe Asp Val Met Gly Tyr Cys Val Thr  
 50 55 60  
 Thr Ala Lys Gly Ala Ala Glu Thr Leu Lys Leu Asn Met Ala Ile Ile  
 65 70 75 80  
 Leu Leu Pro Val Cys Arg Asn Thr Ile Thr Trp Leu Arg Ser Thr Arg  
 85 90 95  
 Ala Ala Arg Ala Leu Pro Phe Asp Asp Asn Ile Asn Phe His Lys Thr  
 100 105 110  
 Ile Ala Ala Ala Ile Val Val Gly Ile Ile Leu His Ala Gly Asn His  
 115 120 125  
 Leu Val Cys Asp Phe Pro Arg Leu Ile Lys Ser Ser Asp Glu Lys Tyr  
 130 135 140  
 Ala Pro Leu Gly Gln Tyr Phe Gly Glu Ile Lys Pro Thr Tyr Phe Thr  
 145 150 155 160  
 Leu Val Lys Gly Val Glu Gly Ile Thr Gly Val Ile Met Val Val Cys  
 165 170 175  
 Met Ile Ile Ala Phe Thr Leu Ala Thr Arg Trp Phe Arg Arg Ser Leu  
 180 185 190  
 Val Lys Leu Pro Arg Pro Phe Asp Lys Leu Thr Gly Phe Asn Ala Phe  
 195 200 205  
 Trp Tyr Ser His His Leu Phe Ile Ile Val Tyr Ile Ala Leu Ile Val  
 210 215 220  
 His Gly Glu Cys Leu Tyr Leu Ile His Val Trp Tyr Arg Arg Thr Thr  
 225 230 235 240  
 Trp Met Tyr Leu Ser Val Pro Val Cys Leu Tyr Val Gly Glu Arg Ile  
 245 250 255



Leu Arg Phe Phe Arg Ser Gly Ser Tyr Ser Val Arg Leu Leu Lys Val  
 260 265 270  
 Ala Ile Tyr Pro Gly Asn Val Leu Thr Leu Gln Met Ser Lys Pro Pro  
 275 280 285  
 Thr Phe Arg Tyr Lys Ser Gly Gln Tyr Met Phe Val Gln Cys Pro Ala  
 290 295 300  
 Val Ser Pro Phe Glu Trp His Pro Phe Ser Ile Thr Ser Ala Pro Gly  
 305 310 315 320  
 Asp Asp Tyr Leu Ser Ile His Val Arg Gln Leu Gly Asp Trp Thr Arg  
 325 330 335  
 Glu Leu Lys Arg Val Phe Ala Ala Ala Cys Glu Pro Pro Ala Gly Gly  
 340 345 350  
 Lys Ser Gly Leu Leu Arg Ala Asp Glu Thr Thr Lys Lys Ile Leu Pro  
 355 360 365  
 Lys Leu Leu Ile Asp Gly Pro Tyr Gly Ser Pro Ala Gln Asp Tyr Ser  
 370 375 380  
 Lys Tyr Asp Val Leu Leu Leu Val Gly Leu Gly Ile Gly Ala Thr Pro  
 385 390 395 400  
 Phe Ile Ser Ile Leu Lys Asp Leu Leu Asn Asn Ile Ile Lys Met Glu  
 405 410 415  
 Glu Glu Glu Asp Ala Ser Thr Asp Leu Tyr Pro Pro Met Gly Arg Asn  
 420 425 430  
 Asn Pro His Val Asp Leu Gly Thr Leu Met Thr Ile Thr Ser Arg Pro  
 435 440 445  
 Lys Lys Ile Leu Lys Thr Thr Asn Ala Tyr Phe Tyr Trp Val Thr Arg  
 450 455 460  
 Glu Gln Gly Ser Phe Asp Trp Phe Lys Gly Val Met Asn Glu Ile Ala  
 465 470 475 480  
 Asp Leu Asp Gln Arg Asn Ile Ile Glu Met His Asn Tyr Leu Thr Ser  
 485 490 495  
 Val Tyr Glu Glu Gly Asp Ala Arg Ser Ala Leu Ile Thr Met Leu Gln  
 500 505 510



Ala Leu Asn His Ala Lys Asn Gly Val Asp Ile Val Ser Gly Thr Lys  
 515 520 525

Val Arg Thr His Phe Ala Arg Pro Asn Trp Arg Lys Val Leu Ser Lys  
 530 535 540

Ile Ser Ser Lys His Pro Tyr Ala Lys Ile Gly Val Phe Tyr Cys Gly  
 545 550 555 560

Ala Pro Val Leu Ala Gln Glu Leu Ser Lys Leu Cys His Glu Phe Asn  
 565 570 575

Gly Lys Cys Thr Thr Lys Phe Asp Phe His Lys Glu His Phe  
 580 585 590

<210> 41

<211> 2619

<212> DNA

<213> Rat

<400> 41

gtgctgtcag agctttacag agcctctggg catgcgcatt gctacccatt tcattgattt 60  
 acagaagtca tgctaaaatc tctttcatgc atgtcttctt ttttcagtct ctccctttccc 120  
 aaagcttttc agtttgcctt ttgcttgtag caactgctat cctcctcaa aggctgctgc 180  
 aaaaggtagt cctttttctt ggaggctttc agcaaatact acctgggaac ctgcttcagc 240  
 tcttggaata ttaagtga gagaacattt catagcattt gtatctttct ttgaaggagc 300  
 caccagacag actgccttgg ccttgccag agcatctgct ttgtgcctga attttaacag 360  
 catggtgatc ctgattcctg tgtgtcgaaa tctgctctcc ttcctgaggg gcacctgctc 420  
 attttgcaac cacacgctga gaaagccatt ggatcacaac ctcaccttcc ataagctggc 480  
 ggcatatatg atctgcatat tcacagctat tcatatcatt gcacatctat ttaactttga 540  
 acgctacagt agaagccaac aggccatgga tggatctctt gcctctgttc tctccagcct 600  
 attccatccc gagaaagaag attcttggct aaatcccatc cagtctccaa acgtgacagt 660  
 gatgtatgca gcatttacca gtattgctgg ccttactgga gtggctcgcca ctgtggcttt 720  
 ggttctcatg gtaacttcag ctatggagtt tatccgcagg aattatcttg agctcttctg 780



gtatacacat caccttttca tcatctatat catctgctta gggatccatg gcctgggggg 840  
gattgtccgg ggtcaaacag aagagagcat gagtgaaagt catccccgca actgttcata 900  
ctctttccac gagtgggata agtatgaaag gagttgcagg agtcctcatt ttgtggggca 960  
acccctgag tcttgggaagt ggatcctcgc gccgattgct ttttatatct ttgaaaggat 1020  
ccttcgcttt tatcgctccc ggcagaaggt cgtgattacc aaggttgca tgcacccatg 1080  
taaagttttg gaattgcaga tgaggaagcg gggctttact atgggaatag gacagtatat 1140  
attcgtaa at tgccccctga tttccttcct ggaatggcat ccctttactc tgacctctgc 1200  
tccagaggaa gaatttttct ccattcatat tcgagcagca ggggactgga cagaaaatct 1260  
cataaggaca tttgaacaac agcactcacc aatgccagg atcgaggtgg atggtccctt 1320  
tggcacagtc agtgaggatg tcttcagta cgaagtggct gtactggttg gggcaggat 1380  
tggcgtcact cccttgctt ccttcttgaa atctatctgg taaaaattcc agcgtgcaca 1440  
caacaagctg aaaacacaaa agatctatct ctactggatt tgtagagaga cgggtgcctt 1500  
tgcttggttc aacaacttat tgaattccct ggaacaagag atggacgaat taggcaaacc 1560  
ggatttccta aactaccgac tcttcctcac tggctgggat agcaacattg ctggatcatgc 1620  
agcattaaac tttgacagag ccactgacgt cctgacaggt ctgaaacaga aaacctcctt 1680  
tgaggagacca atgtgggaca atgagttttc tagaatagct actgccacc ccaagtctgt 1740  
gggtgggggtt ttcttatgcg gccctccgac tttggcaaaa agcctgcgca aatgctgtcg 1800  
gcggtactca agtctggatc ctaggaaggt tcaattctac ttcaacaaag aaacgttctg 1860  
aattggagga agccgcacag tagtacttct ccattctcct tttcactaac gtgtgggtca 1920  
gctactagat agtccgttgt cgcacaagga cttcactccc atcttaaagt tgactcaact 1980  
ccatcattct tgggcttttg caacatgaga gctgcataac tcacaattgc aaaacacatg 2040  
aattattatt ggggggattg taaatccttc tgggaaacct gcctttagct gaatcttgct 2100  
ggttgacact tgcacaattt aacctcaggt gtcttggtg atacctgata atcttcctc 2160  
ccacctgtcc ctcacagaag atttctaagt agggtgattt taaaatattt attgaatcca 2220



cgacaaaaca ataatacataa ataataaaca taaaattacc aagattccca ctcccatatc 2280  
 ataccacta agaacatcgt tatacatgag cttatcatcc agtgtgacca acaatttata 2340  
 ctttactgtg ccaaaataat cttcatcttt gcttattgaa caattttgct gactttccct 2400  
 agtaatatct taagtatatt aactggaatc aaatttgtat tatagttaga agccaactat 2460  
 attgccagtt tgtattgttt gaaataactg gaaaggcctg acctacatcg tggggtaatt 2520  
 taacagaagc tctttccatt ttttgttgtt gttgttaaag agttttgttt atgaatgtgt 2580  
 tataaaaaga aaataaaaag ttataatddd gacggaaaa 2619

&lt;210&gt; 42

&lt;211&gt; 499

&lt;212&gt; PRT

&lt;213&gt; Rat

&lt;400&gt; 42

Met Val Ile Leu Ile Pro Val Cys Arg Asn Leu Leu Ser Phe Leu Arg  
 1 5 10 15

Gly Thr Cys Ser Phe Cys Asn His Thr Leu Arg Lys Pro Leu Asp His  
 20 25 30

Asn Leu Thr Phe His Lys Leu Val Ala Tyr Met Ile Cys Ile Phe Thr  
 35 40 45

Ala Ile His Ile Ile Ala His Leu Phe Asn Phe Glu Arg Tyr Ser Arg  
 50 55 60

Ser Gln Gln Ala Met Asp Gly Ser Leu Ala Ser Val Leu Ser Ser Leu  
 65 70 75 80

Phe His Pro Glu Lys Glu Asp Ser Trp Leu Asn Pro Ile Gln Ser Pro  
 85 90 95

Asn Val Thr Val Met Tyr Ala Ala Phe Thr Ser Ile Ala Gly Leu Thr  
 100 105 110

Gly Val Val Ala Thr Val Ala Leu Val Leu Met Val Thr Ser Ala Met  
 115 120 125

Glu Phe Ile Arg Arg Asn Tyr Phe Glu Leu Phe Trp Tyr Thr His His  
 130 135 140



Leu Phe Ile Ile Tyr Ile Ile Cys Leu Gly Ile His Gly Leu Gly Gly  
 145 150 155 160  
 Ile Val Arg Gly Gln Thr Glu Glu Ser Met Ser Glu Ser His Pro Arg  
 165 170 175  
 Asn Cys Ser Tyr Ser Phe His Glu Trp Asp Lys Tyr Glu Arg Ser Cys  
 180 185 190  
 Arg Ser Pro His Phe Val Gly Gln Pro Pro Glu Ser Trp Lys Trp Ile  
 195 200 205  
 Leu Ala Pro Ile Ala Phe Tyr Ile Phe Glu Arg Ile Leu Arg Phe Tyr  
 210 215 220  
 Arg Ser Arg Gln Lys Val Val Ile Thr Lys Val Val Met His Pro Cys  
 225 230 235 240  
 Lys Val Leu Glu Leu Gln Met Arg Lys Arg Gly Phe Thr Met Gly Ile  
 245 250 255  
 Gly Gln Tyr Ile Phe Val Asn Cys Pro Ser Ile Ser Phe Leu Glu Trp  
 260 265 270  
 His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Glu Phe Phe Ser Ile  
 275 280 285  
 His Ile Arg Ala Ala Gly Asp Trp Thr Glu Asn Leu Ile Arg Thr Phe  
 290 295 300  
 Glu Gln Gln His Ser Pro Met Pro Arg Ile Glu Val Asp Gly Pro Phe  
 305 310 315 320  
 Gly Thr Val Ser Glu Asp Val Phe Gln Tyr Glu Val Ala Val Leu Val  
 325 330 335  
 Gly Ala Gly Ile Gly Val Thr Pro Phe Ala Ser Phe Leu Lys Ser Ile  
 340 345 350  
 Trp Tyr Lys Phe Gln Arg Ala His Asn Lys Leu Lys Thr Gln Lys Ile  
 355 360 365  
 Tyr Phe Tyr Trp Ile Cys Arg Glu Thr Gly Ala Phe Ala Trp Phe Asn  
 370 375 380  
 Asn Leu Leu Asn Ser Leu Glu Gln Glu Met Asp Glu Leu Gly Lys Pro  
 385 390 395 400



Asp Phe Leu Asn Tyr Arg Leu Phe Leu Thr Gly Trp Asp Ser Asn Ile  
 405 410 415

Ala Gly His Ala Ala Leu Asn Phe Asp Arg Ala Thr Asp Val Leu Thr  
 420 425 430

Gly Leu Lys Gln Lys Thr Ser Phe Gly Arg Pro Met Trp Asp Asn Glu  
 435 440 445

Phe Ser Arg Ile Ala Thr Ala His Pro Lys Ser Val Val Gly Val Phe  
 450 455 460

Leu Cys Gly Pro Pro Thr Leu Ala Lys Ser Leu Arg Lys Cys Cys Arg  
 465 470 475 480

Arg Tyr Ser Ser Leu Asp Pro Arg Lys Val Gln Phe Tyr Phe Asn Lys  
 485 490 495

Glu Thr Phe

<210> 43  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: synthetic  
 primer

<400> 43  
 ttctgagtag gtgtgcattt gagtgcata aagac 35

<210> 44  
 <211> 45  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: synthetic  
 primer

<400> 44  
 ttttccgtca aaattataac tttttatttt ctttttataa cacat 45



<210> 45  
<211> 5508  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (155)..(4810)

<400> 45

```
1   GCAGAGCTGC AGAGGCACCG GACGAGAGAG GGCTCCGCGG GCCCAGCTGG CAGCCAGGCC
61  GGAGACAAGT TGCAGTCCCG GGCTCTGGTG ACGCCGTGGC CGCAGGGTCT CCATTTTGGG
121 ACATTCTAAT CCCTGAGCCC CTATTATTTT CATCATGGGC TTCTGCCTGG CTCTAGCATG
181 GACACTTCTG GTTGGGGCAT GACCCCTCT GGGAGCTCAG AACCCCATTT CGTGGGAGGT
241 GCAGCGATTT GATGGGTGGT ACAACAACCT CATGGAGCAC AGATGGGGCA GCAAAGGCTC
301 CCGGCTGCAG CGCCTGGTCC CAGCCAGCTA TGCAGATGGC GTGTACCAGC CCTTGGGAGA
361 ACCCCACCTG CCCAACCCCC GAGACCTTAG CAACACCATC TCAAGGGGCC CTGCAGGGCT
421 GGCCTCCCTG AGAAACCGCA CAGTGTTCGG GGTCTTCTTT GGCTATCACG TGCTTTCAGA
481 CCTGGTGAGC GTGGAAACTC CCGGCTGCCC CGCCGAGTTC CTCAACATTC GCATCCCGCC
541 CGGAGACCCC ATGTTTCGACC CCGACCAGCG CGGGGACGTG GTGCTGCCCT TCCAGAGAAG
601 CCGCTGGGAC CCCGAGACCG GACGGAGTCC CAGCAATCCC CGGGACCCGG CCAACCAGGT
661 GACGGCTGGT CTGGACGGCA GCGCCATCTA TGGTTCCTCG CATTCTTGGA GCGACGCGCT
721 GCGGAGCTTC TCCAGGGGAC AGCTGGCGTC GGGGCCCAGC CCCGCTTTTC CCCGAGACTC
781 GCAGAACCCC CTGCTCATGT GGGCGGCGCC CGACCCCGCC ACCGGGCAGA ACGGGCCCCG
841 GGGGCTGTAC GCCTTCGGGG CAGAGAGAGG GAACCGGGAA CCCTTCCTGC AGGCGCTGGG
901 CCTGCTCTGG TTCCGCTACC ACAACCTGTG GGCGCAGAGG CTGGCCCGCC AGCACCAGAA
961 CTGGGAGGAG GAGGAGCTGT TCCAGCACGC ACGCAAGAGG GTCATCGCCA CCTACCAGAA
1021 CATCGCTGTG TATGAGTGGC TGCCCAGCTT CCTGCAGAAA AACTTCCCGG AGTATACAGG
1081 ATACCGGCCA TTTCTGGACC CCAGCATCTC CTCAGAGTTC GTGGCGGCCCT CTGAGCAGTT
1141 CCTGTCCACC ATGGTGCCCC CTGGCGTCTA CATGAGAAAT GCCAGCTGCC ACTTCCAGGG
1201 GGTCAATCAAT CGGAACCTAA GTGTCTCCAG AGTCTCCGG GTCTGCAACA GCTACTGGAG
1261 CCGTGAGCAC CCAAGCCTAC AAAGTGCTGA AGATGTGGAT GCACTGCTGC TGGGCATGGC
1321 CTCCCAGATC GCAGAGCGAG AGGACCATGT GTTGGTTGAA GATGTGCGGG ATTTCTGGCC
1381 TGGGCCACTG AAGTTTTCCT GCACAGACCA CCTGGCCAGC TGCCTGCGAG GGGGCCGGGA
1441 TCTGGGCCTG CCCTCTTACA CCAAGGCCAG GGCAGCACTG GGCTTGCTC CCATTACCCG
1501 CTGGCAGGAC ATCAACCCCTG CACTCTCCCG GAGCAATGAC ACTGTACTGG AGGCCACAGC
1561 TGCCCTGTAC AACCAGGACT TATCCTGGCT AGAGCTGCTC CCTGGGGGAC TCCTGGAGAG
1621 CCACCGGGAC CCTGGACCTC TGTTACAGC CATCGTCTT GAACAATTTG TGCGGCTACG
1681 GGATGGTGAC CGCTACTGGT TTGAGAACAC CAGGAATGGG CTGTTCTCCA AGAAGGAGAT
1741 TGAAGAAATC CGAAATACCA CCCTGCAGGA CGTGCTGGTC GCTGTTATCA ACATTGACCC
1801 CAGTGCTCTG CAGCCCAATG TCTTTGTCTG GCATAAAGGA GACCCCTGTC CGCAGCCGAG
1861 ACAGCTCAGC ACTGAAGGCC TGCCAGCGTG TGCTCCCTCT GTTGTTCGTG ACTATTTTGA
1921 GGGCAGTGGA TTTGGCTTCG GGGTCACCAT CGGGACCTC TGTGCTTCC CTTTGGTGAG
1981 CCTGCTCAGT GCCTGGATTG TTGCCCAGCT CCGGATGAGA AATTTCAGGA GGCTCCAGGG
2041 CCAGGACCGC CAGAGCATCG TGTCTGAGAA GCTCGTGGGA GGCATGGAAG CTTTGGGAATG
2101 GCAAGGCCAC AAGGAGCCCT GCCGGCCCGT GCTTGTGTAC CTGCAGCCCG GGCAGATCCG
2161 TGTGGTAGAT GGCAGGCTCA CCGTGCTCCG CACCATCCAG CTGCAGCCTC CACAGAAGGT
2221 CAACTTCGTC CTGTCCAGCA ACCGTGGACG CCGCACTCTG CTGCTCAAGA TCCCCAAGGA
2281 GTATGACCTG GTGCTGCTGT TTAAGTTGGA GGAAGAGCGG CAGGCGCTGG TGAAAAATCT
2341 CCGGGGAGCT CTGAAGGAGA GCGGGTTGAG CATCCAGGAG TGGGAGCTGC GGGAGCAGGA
```



2401 GCTGATGAGA GCAGCTGTGA CACGGGAGCA GCGGAGGCAC CTCCTGGAGA CCTTTTTCAG  
 2461 GCACCTTTTC TCCCAGGTGC TGGACATCAA CCAGGCCGAC GCAGGGACCC TCCCCCTGGA  
 2521 CTCCTCCCAG AAGGTGCGGG AGGCCCTGAC CTGTGAGCTG AGCAGGGCCG AGTTTGCCGA  
 2581 GTCCCTGGGC CTCAAGCCCC AGGACATGTT TGTGGAGTCC ATGTTCTCTC TGGCTGACAA  
 2641 GGATGGCAAT GGCTACCTGT CCTTCCGAGA GTTCCCTGGAC ATCCTGGTGG TCTTCATGAA  
 2701 AGGCTCTCCT GAGGAAAAGT CTCGCCTTAT GTTCCGCATG TACGACTTTG ATGGGAATGG  
 2761 CCTCATTTCC AAGGATGAGT TCATCAGGAT GCTGAGATCC TTCATCGAGA TCTCCAACAA  
 2821 CTGCCTGTCC AAGGCCCAGC TGGCTGAGGT GGTGGAGTCC ATGTTCCGGG AGTCGGGATT  
 2881 CCAGGACAAG GAGGAACTGA CATGGGAAGA TTTTCACTTC ATGCTGCGGG ACCACAATAG  
 2941 CGAGCTCCGC TTCACGCAGC TCTGTGTCAA AGGGGTGGAG GTGCCTGAAG TCATCAAGGA  
 3001 CCTCTGCCGG CGAGCCTCCT ACATCAGCCA GGATATGATC TGTCCCTCTC CCAGAGTGAG  
 3061 TGCCCGCTGT TCCCGCAGCG ACATTGAGAC TGAGTTGACA CCTCAGAGAC TGCAGTGCCC  
 3121 CATGGACACA GACCTCCCC AGGAGATTCTG GCGGAGGTTT GGCAAGAAGG TAACGTCATT  
 3181 CCAGCCCTTG CTGTTCACTG AGGCGCACCG AGAGAAGTTC CAACGCAGCT GTCTCCACCA  
 3241 GACGGTGCAA CAGTTCAAGC GCTTCATTGA GAACTACCGG CGCCACATCG GCTGCGTGGC  
 3301 CGTGTCTTAC GCCATCGCTG GGGGGCTTTT CCTGGAGAGG GCCTACTACT ACGCCTTTGC  
 3361 CGCACATCAC ACGGGCATCA CGGACACCAC CCGCGTGGGA ATCATCCTGT GGGGGGGCAC  
 3421 AGCAGCCAGC ATCTCTTTCA TGTCTCCTA CATCTTGCTC ACCATGTGCC GCAACCTCAT  
 3481 CACCTTCCTG CGAGAAACCT TCCTCAACCG CTACGTGCCC TTCGACGCCG CCGTGGACTT  
 3541 CCATCGCCTC ATTGCCTCCA CCGCCATCGT CCTCACAGTC TTACACAGTG TGGGCCATGT  
 3601 GGTGAATGTG TACCTGTTCT CCATCAGCCC CCTCAGCGTC CTCTCTTGCC TCTTTCTGCG  
 3661 CCTCTTCCAT GATGATGGGT CTGAGTTCCC CCAGAAGTAT TACTGGTGGT TCTTCCAGAC  
 3721 CGTACCAGGC CTCACGGGGG TTGTGCTGCT CCTGATCCTG GCCATCATGT ATGCTTTTGC  
 3781 CTCCCACCAC TTCCGCCGCC GCAGTTTCCG GGGCTTCTGG CTGACCCACC ACCTCTACAT  
 3841 CTTGCTCTAT GTCCGTGCTCA TCATCCATGG TAGCTTTGCC CTGATCCAGC TGCCCCGTTT  
 3901 CCACATCTTC TTCTGTGTC CAGCAATCAT CTATGGGGGC GACAAGCTGG TGAGCCTGAG  
 3961 CCGGAAGAAG GTGGAGATCA GCGTGGTGAA GCGGAGCTG CTGCCCTCAG GAGTGACCCA  
 4021 CCTGCGGTTT CAGCGGCCCC AGGGCTTTGA GTACAAGTCA GGGCAGTGGG TGCGGATCGC  
 4081 TTGCCTGGCT CTGGGGACCA CCGAGTACCA CCCCTTCACA CTGACCTCTG CGCCCCATGA  
 4141 GGACACGCTT AGCCTGCACA TCCGGGCAGC AGGGCCCTGG ACCACTCGCC TCAGGGAGAT  
 4201 CTA CTAGCC CCGACGGGTG ACAGATGTGC CAGATACCCA AAGCTGTACC TTGATGGACC  
 4261 ATTTGGAGAG GGCCACCAGG AGTGGCATAA GTTTGAGGTG TCAGTGTTAG TGGGAGGGGG  
 4321 CATTGGGGTC ACCCTTTTGG CCTCCATCCT CAAAGACCTG GTCTTCAAGT CATCCGTCAG  
 4381 CTGCCAAGTG TTCTGTAAAG AGATCTACTT CATCTGGGTG ACGCGGACCC AGCGTCAGTT  
 4441 TGAGTGGCTG GCTGACATCA TCCGAGAGGT GGAGGAGAAT GACCACCAGG ACCTGGTGTC  
 4501 TGTGCACATC TACATCACCC AGCTGGCTGA GAAGTTCGAC CTCAGGACCA CTATGCTGTA  
 4561 CATCTGTGAG CGGCACTTCC AGAAGGTTCT GAACCGGAGT CTATTACAG GCCTGCGCTC  
 4621 CATCACCCAC TTTGGCCGTC CCCCCTTTGA GCCCTTCTTC AACTCCCTGC AGGAGGTCCA  
 4681 CCCCCAGGTC CGGAAGATCG GGGTGTTTAG CTGTGGCCCC CCTGGCATGA CCAAGAATGT  
 4741 GGAAAAGGCC TGTACGCTCA TCAACAGGCA GGACCGGACT CACTTCTCCC ACCATTATGA  
 4801 GAACTTCTAG GCCCCTGCCC GGGGGTTCTG CCCACTGTCC AGTTGAGCAG AGGTTTGAGC  
 4861 CCACACCTCA CCTCTGTTCT TCCTATTTCT GGCTGCCTCA GCCTTCTCTG ATTTCCCACC  
 4921 TCCCAACCTT GTTCCAGGTG GCCATAGTCA GTCACCATGT GTGGGCTCAG GGACCCCCAG  
 4981 GACCAGGATG TGTCTCAGCC TGGAGAAATG GTGGGGGGGC AGTGTCTAGG GACTAGAGTG  
 5041 AGAAGTAGGG GAGCTACTGA TTTGGGGCAA AGTGAAACCT CTGCTTCAGA CTTCAGAAAC  
 5101 AAATCTCAGA AGACAAGCTG ACCTGACAAG TACTATGTGT GTGCATGTCT GTATGTGTGT  
 5161 TGGGGCGGTG AGTGTAAGGA TGCAGTGGGA GCATGGATGC TGGCATCTTA GAACCCCTCC  
 5221 TACTCCATA CCTCTCCTC TTCTGGGCTC CCCACTGTCA GACGGGCTGG CAAATGCCTT  
 5281 GCAGGAGGTA GAGGCTGGAC CCATGGCAAG CCATTTACAG AAACCCACTC GGCACCCAG



5341 TCTAACACCA CAACTAATT CACCCAAGGT TTTAAGCACG TTCTTTCATC AGACCCTGGC  
5401 CCAATACCTA TGTATGCAAT GTCCTCAGC CCTCTTCTCC CTGCTCCAGT AGTCTCCCTT  
5461 CCAAATAAAT CACTTTTCTG CCAAAAAAAAA AAAA



<210> 46  
<211> 1521  
<212> PRT  
<213> Homo sapiens  
  
<400> 46



1 MGFCLALAWT LLVGAWTPLG AQNPISWEVQ RFDGWYNNLM EHRWGSKGSR LQRLVPASYA  
 61 DGVYQPLGEP HLPNPRDLSN TISRGPAGLA SLRNRTVLGV FFGYHVLSDL VSVETPGCPA  
 121 EFLNIRIPPG DPMFDPDQRG DVVLPPQRSR WDPETGRSPS NPRDPANQVT GWLDGSAIYG  
 181 SSHSWSDALR SFSRQQLASG PDPAFFRDSQ NPLLMWAAPD PATGQNGPRG LYAFGAERGN  
 241 REPFLQALGL LWFRYHNLWA QRLARQHPDW EDEELFQHAR KRVIATYQNI AVYEWLPSFL  
 301 QKTLPEYTG YRPFLDPSISS EFVAASEQFL STMVPPGVYM RNASCHFQGV INRNSSVSRA  
 361 LRVCSYWSR EHPSLQSAED VDALLGMSA QIAEREDHVL VEDVRDFWPG PLKFSRTDHL  
 421 ASCLQRGRDL GLPSYTKARA ALGLSPITRW QDINPALSRS NDTVLEATAA LYNQDLSWLE  
 481 LLPGGLLSH RDGGLPSTI VLEQFVRLRD GDRYWFENTR NGLFSKKEIE EIRNTTLQDV  
 541 LVAVINIDPS ALQPNVFWH KGDPCPQPRQ LSTEGLPACA PSVVRDYFEG SGFGFGVTIG  
 601 TLCCFPLVSL LSAWIVARLR MRNFKRLQGG DRQSIVSEKL VGGMEALEWQ GHKEPCRPVL  
 661 VYLQPGQIRV VDGRITVLRT IQLQPPQKVN FVLSSNRGRR TLLKIPKEY DLVLLFNLEE  
 721 ERQALVENLR GALKESGLSI QEWELEQEL MRAAVTREQR RHLLETFFRH LFSQVLDINQ  
 781 ADAGTLPLDS SQKVRALTC ELSRAEFAES LGLKPDQMFV ESMFSLADKD GNGYLSFREF  
 841 LDILVVFMMG SPEEKSRMF RMYDFDGNGL ISKDEFIRML RSFIEISNNC LSQAQLAEV  
 901 ESMFRESGQ DKEELTWEDF HFMLRDHNS LRFTQLCVKG VEVPEVIKDL CRRASYISQD  
 961 MICSPRVSAR CSRSRDIETE LTPQRLQCPM DTDPPQEIRR RFGKKVTSFQ PLLFTEAHRE  
 1021 KFORSLHQT VQFQKRFIEN YRRHIGCVAV FYAIAGGLFL ERAYYYAFAA HHTGITDTTR  
 1081 VGIILSRGTA ASISFMFSYI LLTMCRLIT FLRETFLNRY VPFDAVDFH RLIATAIVL  
 1141 TVLHVSQHV NVYLFSSPL SVLSCLFPGL FHDDGSEFPQ KYWVWFQTV PGLTGVVLLL  
 1201 ILAIMYVFAS HHFRRSFRG FWLTHHLYIL LYVLLIINGS FALIQLPRFH IFFLVPAIY  
 1261 GGDKLVSLSR KKVEISVKA ELLPSGVTHL RFQRPQGFY KSGQWVRIAC LALGTTEYHP  
 1321 FTLTSAPHED TSLSHIRAAG PWTTRLREIY SAPTGDRCAR YPKLYLDGPF GEGHQEWHKF  
 1381 EVSVLVGGGI GVTFFASILK DLVFKSSVSC QVFCKKIYFI WVTRTQRQFE WLADIIREVE  
 1441 ENDHQDLVSV HIYITQLAEK FDLRTTMLYI CERHFQKVLN RSLFTGLRSI THFGRPPFEP  
 1501 FFNSLQEVHP QVRKIGVFSC GPPGMTKNVE KACQLINRQD RTHFSHHYEN F.



WO 00/28031

PCT/US99/26592

<210> 47



&lt;211&gt; 3453

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (438)..(3134)

&lt;400&gt; 47

```

gtcctcgacc agtttgtacg gctgcgggat ggtgaccgct actggtttga gaacaccagg 60
aatgggctgt tctccaagaa ggagattgag acatccgaaa taccaccgtg cgggacgtgc 120
tggtcgctgt tatcaacatt gacccacgtg ccctgcagcc caatgtcttt gtctggcata 180
aaggtgcacc ctgccctcaa cctaagcagc tcacaactga cggcctgccc cagtgtgcac 240
ccctgactgt gcttgacttc tttgaaggca gcagccctgg ttttgccatc accatcattg 300
ctctctgctg ccttccctta gtgagtctgc ttctctctgg agtgggtggc tatttcggg 360
gccgagaaca caagaagcta caaaagaaac tcaaagagag cgtgaagaag gaagcagcca 420
aagatggagt gccagcg atg gag tgg cca ggc ccc aag gag agg agc agt      470
                Met Glu Trp Pro Gly Pro Lys Glu Arg Ser Ser
                  1              5                  10
ccc atc atc atc cag ctg ctg tca gac agg tgt ctg cag gtc ctg aac      518
Pro Ile Ile Ile Gln Leu Leu Ser Asp Arg Cys Leu Gln Val Leu Asn
          15              20                  25
agg cat ctc act gtg ctc cgt gtg gtc cag ctg cag cct ctg cag cag      566
Arg His Leu Thr Val Leu Arg Val Val Gln Leu Gln Pro Leu Gln Gln
          30              35                  40
gtc aac ctc atc ctg tcc aac aac cga gga tgc cgc acc ctg ctg ctc      614
Val Asn Leu Ile Leu Ser Asn Asn Arg Gly Cys Arg Thr Leu Leu Leu
          45              50                  55
aag atc cct aag gag tat gac ctg gtg ctg ctg ttt agt tct gaa gag      662
Lys Ile Pro Lys Glu Tyr Asp Leu Val Leu Leu Phe Ser Ser Glu Glu
          60              65              70              75
gaa cgg ggc gcc ttt gtg cag cag cta tgg gac ttc tgc gtg cgc tgg      710
Glu Arg Gly Ala Phe Val Gln Gln Leu Trp Asp Phe Cys Val Arg Trp
          80              85              90
gct ctg ggc ctc cat gtg gct gag atg agc gag aag gag cta ttt agg      758

```



Ala Leu Gly Leu His Val Ala Glu Met Ser Glu Lys Glu Leu Phe Arg	
95 100 105	
aag gct gtg aca aag cag cag cgg gaa cgc atc ctg gag atc ttc ttc	806
Lys Ala Val Thr Lys Gln Gln Arg Glu Arg Ile Leu Glu Ile Phe Phe	
110 115 120	
aga cac ctt ttt gct cag gtg ctg gac atc aac cag gcc gac gca ggg	854
Arg His Leu Phe Ala Gln Val Leu Asp Ile Asn Gln Ala Asp Ala Gly	
125 130 135	
acc ctg ccc ctg gac tcc tcc cag aag gtg cgg gag gcc ctg acc tgc	902
Thr Leu Pro Leu Asp Ser Ser Gln Lys Val Arg Glu Ala Leu Thr Cys	
140 145 150 155	
gag ctg agc agg gcc gag ttt gcc gag tcc ctg ggc ctc aag ccc cag	950
Glu Leu Ser Arg Ala Glu Phe Ala Glu Ser Leu Gly Leu Lys Pro Gln	
160 165 170	
gac atg ttt gtg gag tcc atg ttc tct ctg gct gac aag gat ggc aat	998
Asp Met Phe Val Glu Ser Met Phe Ser Leu Ala Asp Lys Asp Gly Asn	
175 180 185	
ggc tac ctg tcc ttc cga gag ttc ctg gac atc ctg gtg gtc ttc atg	1046
Gly Tyr Leu Ser Phe Arg Glu Phe Leu Asp Ile Leu Val Val Phe Met	
190 195 200	
aaa ggc tcc cca gag gat aag tcc cgt cta atg ttt acc atg tat gac	1094
Lys Gly Ser Pro Glu Asp Lys Ser Arg Leu Met Phe Thr Met Tyr Asp	
205 210 215	
ctg gat gag aat ggc ttc ctc tcc aag gac gaa ttc ttc acc atg atg	1142
Leu Asp Glu Asn Gly Phe Leu Ser Lys Asp Glu Phe Phe Thr Met Met	
220 225 230 235	
cga tcc ttc atc gag atc tcc aac aac tgc ctg tcc aag gcc cag ctg	1190
Arg Ser Phe Ile Glu Ile Ser Asn Asn Cys Leu Ser Lys Ala Gln Leu	
240 245 250	
gcc gag gtg gtg gag tct atg ttc cgg gag tcg gga ttc cag gac aag	1238
Ala Glu Val Val Glu Ser Met Phe Arg Glu Ser Gly Phe Gln Asp Lys	
255 260 265	
gag gag ctg aca tgg gag gat ttt cac ttc atg ctg cgg gac cat gac	1286
Glu Glu Leu Thr Trp Glu Asp Phe His Phe Met Leu Arg Asp His Asp	
270 275 280	
agc gag ctc cgc ttc acg cag ctc tgt gtc aaa ggt gga ggt gga ggt	1334



Ser Glu Leu Arg Phe Thr Gln Leu Cys Val Lys Gly Gly Gly Gly Gly	
285	290 295
gga aat ggt att aga gat atc ttt aaa caa aac atc agc tgt cga gtc	1382
Gly Asn Gly Ile Arg Asp Ile Phe Lys Gln Asn Ile Ser Cys Arg Val	
300	305 310 315
tcg ttc atc act cgg aca cct ggg gag cgc tcc cac ccc cag gga ctg	1430
Ser Phe Ile Thr Arg Thr Pro Gly Glu Arg Ser His Pro Gln Gly Leu	
	320 325 330
ggg ccc cct gtc cca gaa gcc cca gag ctg gga ggc cct gga ctg aag	1478
Gly Pro Pro Val Pro Glu Ala Pro Glu Leu Gly Gly Pro Gly Leu Lys	
	335 340 345
aag agg ttt ggc aaa aag gca gca gtg ccc act ccc cgg ctg tac aca	1526
Lys Arg Phe Gly Lys Lys Ala Ala Val Pro Thr Pro Arg Leu Tyr Thr	
	350 355 360
gag gcg ctg caa gag aag atg cag cga ggc ttc cta gcc caa aag ctg	1574
Glu Ala Leu Gln Glu Lys Met Gln Arg Gly Phe Leu Ala Gln Lys Leu	
	365 370 375
cag cag tac aag cgc ttc gtg gag aac tac cgg agg cac atc gtg tgt	1622
Gln Gln Tyr Lys Arg Phe Val Glu Asn Tyr Arg Arg His Ile Val Cys	
	380 385 390 395
gtg gca atc ttc tcg gcc atc tgt gtt ggc gtg ttt gca gat cgt gct	1670
Val Ala Ile Phe Ser Ala Ile Cys Val Gly Val Phe Ala Asp Arg Ala	
	400 405 410
tac tac tat ggc ttt gcc ttg cca ccc tcg gac att gca cag acc acc	1718
Tyr Tyr Tyr Gly Phe Ala Leu Pro Pro Ser Asp Ile Ala Gln Thr Thr	
	415 420 425
ctc gtg ggc atc atc ctg tca cga ggc acg gcg gcc agc gtc tcc ttc	1766
Leu Val Gly Ile Ile Leu Ser Arg Gly Thr Ala Ala Ser Val Ser Phe	
	430 435 440
atg ttc tct tat atc ttg ctc acc atg tgc cgc aac ctc ata acc ttc	1814
Met Phe Ser Tyr Ile Leu Leu Thr Met Cys Arg Asn Leu Ile Thr Phe	
	445 450 455
ctg cga gag act ttc ctc aac cgc tat gtg cct ttt gat gcc gca gtg	1862
Leu Arg Glu Thr Phe Leu Asn Arg Tyr Val Pro Phe Asp Ala Ala Val	
	460 465 470 475
gac ttc cac cgc tgg atc gcc atg gct gct gtt gtc ctg gcc att ttg	1910



Asp Phe His Arg Trp Ile Ala Met Ala Ala Val Val Leu Ala Ile Leu	
480 485 490	
cac agt gct ggc cac gca gtc aat gtc tac atc ttc tca gtc agc cca	1958
His Ser Ala Gly His Ala Val Asn Val Tyr Ile Phe Ser Val Ser Pro	
495 500 505	
ctc agc ctg ctg gcc tgc ata ttc ccc aac gtc ttt gtg aat gat ggg	2006
Leu Ser Leu Leu Ala Cys Ile Phe Pro Asn Val Phe Val Asn Asp Gly	
510 515 520	
tcc aag ctt ccc cag aag ttc tat tgg tgg ttc ttc cag acc gtc cca	2054
Ser Lys Leu Pro Gln Lys Phe Tyr Trp Trp Phe Phe Gln Thr Val Pro	
525 530 535	
ggg atg aca ggt gtg ctt ctg ctc ctg gtc ctg gcc atc atg tat gtc	2102
Gly Met Thr Gly Val Leu Leu Leu Leu Val Leu Ala Ile Met Tyr Val	
540 545 550 555	
ttc gcc tcc cac cac ttc cgc cgc cgc agc ttc cgg ggc ttc tgg ctg	2150
Phe Ala Ser His His Phe Arg Arg Arg Ser Phe Arg Gly Phe Trp Leu	
560 565 570	
acc cac cac ctc tac atc ctg ctc tat gcc ctg ctc atc atc cat ggc	2198
Thr His His Leu Tyr Ile Leu Leu Tyr Ala Leu Leu Ile Ile His Gly	
575 580 585	
agc tat gct ctg atc cag ctg ccc act ttc cac atc tac ttc ctg gtc	2246
Ser Tyr Ala Leu Ile Gln Leu Pro Thr Phe His Ile Tyr Phe Leu Val	
590 595 600	
ccg gca atc atc tat gga ggt gac aag ctg gtg agc ctg agc cgg aag	2294
Pro Ala Ile Ile Tyr Gly Gly Asp Lys Leu Val Ser Leu Ser Arg Lys	
605 610 615	
aag gtg gag atc agc gtg gtg aag gcg gag ctg ctg ccc tca gga gtg	2342
Lys Val Glu Ile Ser Val Val Lys Ala Glu Leu Leu Pro Ser Gly Val	
620 625 630 635	
acc tac ctg caa ttc cag agg ccc caa ggc ttt gag tac aag tca gga	2390
Thr Tyr Leu Gln Phe Gln Arg Pro Gln Gly Phe Glu Tyr Lys Ser Gly	
640 645 650	
cag tgg gtg cgg atc gcc tgc ctg gct ctg ggg acc acc gag tac cac	2438
Gln Trp Val Arg Ile Ala Cys Leu Ala Leu Gly Thr Thr Glu Tyr His	
655 660 665	
ccc ttc aca ctg acc tcc gcg ccc cat gag gac aca ctc agc ctg cac	2486



Pro Phe Thr Leu Thr Ser Ala	Pro His Glu Asp Thr Leu Ser Leu His	
670	675	680
atc cgg gca gtg ggg ccc tgg acc act cgc ctc agg gag atc tac tca		2534
Ile Arg Ala Val Gly Pro Trp Thr Thr Arg Leu Arg Glu Ile Tyr Ser		
685	690	695
tcc cca aag ggc aat ggc tgt gct gga tac cca aag ctg tac ctt gat		2582
Ser Pro Lys Gly Asn Gly Cys Ala Gly Tyr Pro Lys Leu Tyr Leu Asp		
700	705	710 715
gga ccg ttt gga gag ggc cat cag gag tgg cat aaa ttt gag gtg tca		2630
Gly Pro Phe Gly Glu Gly His Gln Glu Trp His Lys Phe Glu Val Ser		
	720	725 730
gtg ttg gtg gga ggg ggc att ggg gtc acc ccc ttt gcc tcc atc ctc		2678
Val Leu Val Gly Gly Gly Ile Gly Val Thr Pro Phe Ala Ser Ile Leu		
	735	740 745
aaa gac ctg gtc ttc aag tca tcc ttg ggc agc caa atg ctg tgt aag		2726
Lys Asp Leu Val Phe Lys Ser Ser Leu Gly Ser Gln Met Leu Cys Lys		
	750	755 760
aag atc tac ttc atc tgg gtg aca cgg acc cag cgt cag ttt gag tgg		2774
Lys Ile Tyr Phe Ile Trp Val Thr Arg Thr Gln Arg Gln Phe Glu Trp		
	765	770 775
ctg gct gac atc atc caa gag gtg gag gag aac gac cac cag gac ctg		2822
Leu Ala Asp Ile Ile Gln Glu Val Glu Glu Asn Asp His Gln Asp Leu		
	780	785 790 795
gtg tct gtg cac att tat gtc acc cag ctg gct gag aag ttc gac ctc		2870
Val Ser Val His Ile Tyr Val Thr Gln Leu Ala Glu Lys Phe Asp Leu		
	800	805 810
agg acc acc atg cta tac atc tgc gag cgg cac ttc cag aaa gtg ctg		2918
Arg Thr Thr Met Leu Tyr Ile Cys Glu Arg His Phe Gln Lys Val Leu		
	815	820 825
aac cgg agt ctg ttc acg ggc ctg cgc tcc atc acc cac ttt ggc cgt		2966
Asn Arg Ser Leu Phe Thr Gly Leu Arg Ser Ile Thr His Phe Gly Arg		
	830	835 840
ccc ccc ttc gag ccc ttc ttc aac tcc ctg cag gag gtc cac cca cag		3014
Pro Pro Phe Glu Pro Phe Phe Asn Ser Leu Gln Glu Val His Pro Gln		
	845	850 855
gtg cgc aag atc ggg gtg ttc agc tgc ggc cct cca gga atg acc aag		3062



Val Arg Lys Ile Gly Val Phe Ser Cys Gly Pro Pro Gly Met Thr Lys  
 860 865 870 875

aat gta gag aag gcc tgt cag ctc gtc aac agg cag gac cga gcc cac 3110  
 Asn Val Glu Lys Ala Cys Gln Leu Val Asn Arg Gln Asp Arg Ala His  
 880 885 890

ttc atg cac cac tat gag aac ttc tgagcctgtc ctccctggct gctgcttcca 3164  
 Phe Met His His Tyr Glu Asn Phe  
 895

gtatcctgcc ttctcttctg tgcacctaag ttgccagcc ctgctggcaa tctctccatc 3224

agaatccacc ttaggcctca gctggagggc tgcagagccc ctcccaatat tgggagaata 3284

ttgaccaga caattataca aatgagaaaa ggcattaaaa tttacgtttc tgatgatggc 3344

aaagctcatt ttctattag taactctgct gaagatccat ttattgcaat tcatgctgaa 3404

tctaaattgt aaaatttaaa attaaatgca tgtcctcaaa aaaaaaaaaa 3453

<210> 48  
 <211> 899  
 <212> PRT  
 <213> Homo sapiens

<400> 48  
 Met Glu Trp Pro Gly Pro Lys Glu Arg Ser Ser Pro Ile Ile Ile Gln  
 1 5 10 15

Leu Leu Ser Asp Arg Cys Leu Gln Val Leu Asn Arg His Leu Thr Val  
 20 25 30

Leu Arg Val Val Gln Leu Gln Pro Leu Gln Gln Val Asn Leu Ile Leu  
 35 40 45

Ser Asn Asn Arg Gly Cys Arg Thr Leu Leu Leu Lys Ile Pro Lys Glu  
 50 55 60

Tyr Asp Leu Val Leu Leu Phe Ser Ser Glu Glu Glu Arg Gly Ala Phe  
 65 70 75 80

Val Gln Gln Leu Trp Asp Phe Cys Val Arg Trp Ala Leu Gly Leu His  
 85 90 95

Val Ala Glu Met Ser Glu Lys Glu Leu Phe Arg Lys Ala Val Thr Lys  
 100 105 110



Gln Gln Arg Glu Arg Ile Leu Glu Ile Phe Phe Arg His Leu Phe Ala  
 115 120 125  
 Gln Val Leu Asp Ile Asn Gln Ala Asp Ala Gly Thr Leu Pro Leu Asp  
 130 135 140  
 Ser Ser Gln Lys Val Arg Glu Ala Leu Thr Cys Glu Leu Ser Arg Ala  
 145 150 155 160  
 Glu Phe Ala Glu Ser Leu Gly Leu Lys Pro Gln Asp Met Phe Val Glu  
 165 170 175  
 Ser Met Phe Ser Leu Ala Asp Lys Asp Gly Asn Gly Tyr Leu Ser Phe  
 180 185 190  
 Arg Glu Phe Leu Asp Ile Leu Val Val Phe Met Lys Gly Ser Pro Glu  
 195 200 205  
 Asp Lys Ser Arg Leu Met Phe Thr Met Tyr Asp Leu Asp Glu Asn Gly  
 210 215 220  
 Phe Leu Ser Lys Asp Glu Phe Phe Thr Met Met Arg Ser Phe Ile Glu  
 225 230 235 240  
 Ile Ser Asn Asn Cys Leu Ser Lys Ala Gln Leu Ala Glu Val Val Glu  
 245 250 255  
 Ser Met Phe Arg Glu Ser Gly Phe Gln Asp Lys Glu Glu Leu Thr Trp  
 260 265 270  
 Glu Asp Phe His Phe Met Leu Arg Asp His Asp Ser Glu Leu Arg Phe  
 275 280 285  
 Thr Gln Leu Cys Val Lys Gly Gly Gly Gly Gly Gly Asn Gly Ile Arg  
 290 295 300  
 Asp Ile Phe Lys Gln Asn Ile Ser Cys Arg Val Ser Phe Ile Thr Arg  
 305 310 315 320  
 Thr Pro Gly Glu Arg Ser His Pro Gln Gly Leu Gly Pro Pro Val Pro  
 325 330 335  
 Glu Ala Pro Glu Leu Gly Gly Pro Gly Leu Lys Lys Arg Phe Gly Lys  
 340 345 350  
 Lys Ala Ala Val Pro Thr Pro Arg Leu Tyr Thr Glu Ala Leu Gln Glu  
 355 360 365



Lys Met Gln Arg Gly Phe Leu Ala Gln Lys Leu Gln Gln Tyr Lys Arg  
 370 375 380  
 Phe Val Glu Asn Tyr Arg Arg His Ile Val Cys Val Ala Ile Phe Ser  
 385 390 395 400  
 Ala Ile Cys Val Gly Val Phe Ala Asp Arg Ala Tyr Tyr Tyr Gly Phe  
 405 410 415  
 Ala Leu Pro Pro Ser Asp Ile Ala Gln Thr Thr Leu Val Gly Ile Ile  
 420 425 430  
 Leu Ser Arg Gly Thr Ala Ala Ser Val Ser Phe Met Phe Ser Tyr Ile  
 435 440 445  
 Leu Leu Thr Met Cys Arg Asn Leu Ile Thr Phe Leu Arg Glu Thr Phe  
 450 455 460  
 Leu Asn Arg Tyr Val Pro Phe Asp Ala Ala Val Asp Phe His Arg Trp  
 465 470 475 480  
 Ile Ala Met Ala Ala Val Val Leu Ala Ile Leu His Ser Ala Gly His  
 485 490 495  
 Ala Val Asn Val Tyr Ile Phe Ser Val Ser Pro Leu Ser Leu Leu Ala  
 500 505 510  
 Cys Ile Phe Pro Asn Val Phe Val Asn Asp Gly Ser Lys Leu Pro Gln  
 515 520 525  
 Lys Phe Tyr Trp Trp Phe Phe Gln Thr Val Pro Gly Met Thr Gly Val  
 530 535 540  
 Leu Leu Leu Leu Val Leu Ala Ile Met Tyr Val Phe Ala Ser His His  
 545 550 555 560  
 Phe Arg Arg Arg Ser Phe Arg Gly Phe Trp Leu Thr His His Leu Tyr  
 565 570 575  
 Ile Leu Leu Tyr Ala Leu Leu Ile Ile His Gly Ser Tyr Ala Leu Ile  
 580 585 590  
 Gln Leu Pro Thr Phe His Ile Tyr Phe Leu Val Pro Ala Ile Ile Tyr  
 595 600 605  
 Gly Gly Asp Lys Leu Val Ser Leu Ser Arg Lys Lys Val Glu Ile Ser  
 610 615 620



Val Val Lys Ala Glu Leu Leu Pro Ser Gly Val Thr Tyr Leu Gln Phe  
 625 630 635 640  
 Gln Arg Pro Gln Gly Phe Glu Tyr Lys Ser Gly Gln Trp Val Arg Ile  
 645 650 655  
 Ala Cys Leu Ala Leu Gly Thr Thr Glu Tyr His Pro Phe Thr Leu Thr  
 660 665 670  
 Ser Ala Pro His Glu Asp Thr Leu Ser Leu His Ile Arg Ala Val Gly  
 675 680 685  
 Pro Trp Thr Thr Arg Leu Arg Glu Ile Tyr Ser Ser Pro Lys Gly Asn  
 690 695 700  
 Gly Cys Ala Gly Tyr Pro Lys Leu Tyr Leu Asp Gly Pro Phe Gly Glu  
 705 710 715 720  
 Gly His Gln Glu Trp His Lys Phe Glu Val Ser Val Leu Val Gly Gly  
 725 730 735  
 Gly Ile Gly Val Thr Pro Phe Ala Ser Ile Leu Lys Asp Leu Val Phe  
 740 745 750  
 Lys Ser Ser Leu Gly Ser Gln Met Leu Cys Lys Lys Ile Tyr Phe Ile  
 755 760 765  
 Trp Val Thr Arg Thr Gln Arg Gln Phe Glu Trp Leu Ala Asp Ile Ile  
 770 775 780  
 Gln Glu Val Glu Glu Asn Asp His Gln Asp Leu Val Ser Val His Ile  
 785 790 795 800  
 Tyr Val Thr Gln Leu Ala Glu Lys Phe Asp Leu Arg Thr Thr Met Leu  
 805 810 815  
 Tyr Ile Cys Glu Arg His Phe Gln Lys Val Leu Asn Arg Ser Leu Phe  
 820 825 830  
 Thr Gly Leu Arg Ser Ile Thr His Phe Gly Arg Pro Pro Phe Glu Pro  
 835 840 845  
 Phe Phe Asn Ser Leu Gln Glu Val His Pro Gln Val Arg Lys Ile Gly  
 850 855 860  
 Val Phe Ser Cys Gly Pro Pro Gly Met Thr Lys Asn Val Glu Lys Ala  
 865 870 875 880



Cys Gln Leu Val Asn Arg Gln Asp Arg Ala His Phe Met His His Tyr  
                   885                                  890                                  895

Glu Asn Phe

<210> 49

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 49

cctgacagat gtatttcact acccgag

26

<210> 50

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 50

ggatcggagt cactcccttc gctg

24

<210> 51

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 51

ctagaagctc tccttggtgt aataga

26

<210> 52

<211> 24

<212> DNA

<213> Artificial Sequence



&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 52

atgaacacct ctggggtcag ctga

24

&lt;210&gt; 53

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 53

atgaacacct ctggggtcag ctga

24

&lt;210&gt; 54

&lt;211&gt; 25

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 54

gtcctctgca gcattgttcc tctta

25

&lt;210&gt; 55

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 55

cctgacagat gtatttcact acccag

26

&lt;210&gt; 56

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence



&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 56

ggatcggagt cactcccttc gctg

24

&lt;210&gt; 57

&lt;211&gt; 25

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 57

aatgacactg tactggaggc cacag

25

&lt;210&gt; 58

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 58

ctgccatcta ccacacggat ctgc

24

&lt;210&gt; 59

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 59

cttgccattc caaagcttcc atgc

24

&lt;210&gt; 60

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence



&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 60

gtacaagtca ggacagtggg tgcg

24

&lt;210&gt; 61

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer

&lt;400&gt; 61

tggatgatgt cagccagcca ctca

24